

THE OFFSET AND HOST LIGHT DISTRIBUTIONS OF LONG GAMMA-RAY BURSTS: A NEW VIEW FROM *HST* OBSERVATIONS OF *SWIFT* BURSTSPETER K. BLANCHARD¹, EDO BERGER¹, AND WEN-FAI FONG^{2,3}

Draft version September 29, 2015

ABSTRACT

We present the results of an extensive *Hubble Space Telescope* (*HST*) imaging study of ~ 100 *Swift* long-duration gamma-ray bursts (LGRBs) spanning $0.03 \lesssim z \lesssim 9.4$ using relative astrometry from ground- and space-based afterglow observations to locate the bursts within their host galaxies. Using these data, we measure the distribution of LGRB offsets from their host centers, as well as their relation to the underlying host light distribution. We find that the host-normalized offsets of LGRBs are more centrally concentrated than expected for an exponential disk profile, $\langle R/R_h \rangle = 0.67$, and in particular they are more concentrated than the underlying surface brightness profiles of their host galaxies. The distribution of offsets is inconsistent with the distribution for Type II supernovae (SNe) but consistent with the distribution for Type Ib/c SNe. The fractional flux distribution, with a median value of 0.75, indicates that LGRBs prefer some of the brightest locations in their host galaxies but are not as strongly correlated as previous studies indicated. More importantly, we find a clear correlation between the offset and fractional flux, where bursts at offsets $R/R_h \lesssim 0.5$ exclusively occur at fractional fluxes $\gtrsim 0.6$ while bursts at $R/R_h \gtrsim 0.5$ uniformly trace the light of their hosts. This indicates that the spatial correlation of LGRB locations with bright star forming regions seen in the full sample is dominated by the contribution from bursts at small offset and that LGRBs in the outer parts of galaxies show no preference for unusually bright star forming regions. Finally, we find no evidence for evolution from $z \lesssim 1$ to $z \sim 3$ in the offset or fractional flux distributions. We conclude that LGRBs strongly prefer the bright, inner regions of their hosts indicating that the star formation taking place there is more favorable for LGRB progenitor production. This indicates that another environmental factor beyond metallicity, such as binary interactions or IMF differences, may be operating in the central regions of LGRB hosts.

Subject headings: gamma-ray burst: general

1. INTRODUCTION

Long-duration gamma-ray bursts (LGRBs) are the most energetic explosions known in the universe, with a volumetric rate of $\lesssim 1\%$ of the core-collapse supernova rate. A wide range of studies of their accompanying afterglow emission and host galaxies have been used to shed light on the properties of the bursts and their progenitors. In particular, the association with broad-lined Type Ic supernovae (Type Ic-BL SNe) and exclusive locations in star forming galaxies firmly established that LGRBs result from the deaths of massive stars (e.g., Christensen et al. 2004; Woosley & Bloom 2006; Wainwright et al. 2007). Detailed observations of the host galaxies have also indicated a preference for low metallicity environments (Stanek et al. 2006; Levesque et al. 2010b; Graham & Fruchter 2013), although some bursts occur in environments with solar metallicity (Levesque et al. 2010a,b; Levesque 2012). Similarly, afterglow observations established that LGRBs are powered by relativistic jets with an energy scale of $\sim 10^{51}$ erg (e.g., Frail et al. 2001). These observations support the idea that the progenitors of LGRBs are rapidly-rotating massive stars that undergo core-collapse to form a hyper-accreting black hole (collapsars; Woosley 1993; MacFadyen & Woosley 1999). However, it is unclear at the present whether these massive stars are single or whether they

are subject to a wide range of possible binary interaction scenarios (Podsiadlowski et al. 2004; van den Heuvel & Yoon 2007).

Since direct observations of LGRB progenitors are unlikely due to their low volumetric rate, insight into the nature of the progenitors has to rely on studies of the environments in which LGRBs occur. On a large scale, LGRB hosts have been found to generally be blue, compact, low luminosity, low mass, low metallicity galaxies with high specific star formation rates (e.g., Le Floc'h et al. 2003; Christensen et al. 2004; Wainwright et al. 2007; Savaglio et al. 2009), especially when compared to core-collapse SN hosts (Svensson et al. 2010). These results have led to the notion of a preference for low metallicity progenitors.

The sub-galactic environments of LGRBs can provide additional clues about the nature of the progenitors. To date, two major approaches have been employed in the literature: (i) measuring the offset of LGRBs relative to the centers of their hosts, and (ii) measuring the fractional brightness at the LGRB positions relative to the overall distribution of light within their hosts. In this context, Bloom et al. (2002) combined ground-based afterglow observations and *Hubble Space Telescope* (*HST*) host observations to measure the offsets of 20 LGRBs, and found that they were consistent with being drawn from an exponential disk profile. This led them to conclude that LGRBs were associated with star formation, and to rule out the compact object merger progenitor scenario. However, the small sample size and the large uncertainties in many of the measured offsets prevented a more detailed analysis. Fruchter et al. (2006), on the other hand, argued that the ir-

pblanchard@cfa.harvard.edu

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA² Einstein Fellow³ Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA

regular morphologies of at least some LGRB hosts necessitate the use of the fractional brightness technique (hereafter, fractional flux). Using this analysis for 30 LGRB hosts with *HST* observations, in comparison to 16 core-collapse SNe from the GOODS survey, these authors found that LGRBs are more highly concentrated in the brightest regions of their hosts. Svensson et al. (2010) expanded the sample and reached a similar conclusion, including that LGRBs occur in regions with a higher surface luminosity compared to core-collapse SNe. These results are in tension with the conclusion of Bloom et al. (2002) that LGRBs track an exponential light distribution.

In addition to the tension between the results of previous studies, and the relatively small samples used in these studies, it is important to note other potential challenges. First, the aforementioned studies were based on LGRB samples collected from multiple satellites using different trigger criteria, potentially introducing selection effects that are difficult to quantify. Second, the redshifts were mainly obtained from direct host galaxy spectroscopy (emission lines), leading to a bias toward lower redshifts and more luminous hosts; in conjunction with the small samples this also limited the ability to probe any redshift evolution in the LGRB locations. Finally, these studies did not include “dark” bursts (i.e., bursts that suffer from large rest-frame extinction), leading to a potential bias against dusty environments; at least on the global scale the hosts of dark bursts appear to be more luminous and massive than the general LGRB host sample (Krühler et al. 2011; Perley et al. 2013). These shortcomings can now be overcome with the much larger and uniform sample of *Swift* LGRBs.

Beyond the direct study of LGRB progenitors, the fractional flux distribution of LGRBs has also been used as a point of comparison with core-collapse SNe. Fruchter et al. (2006) found that the core-collapse SNe from GOODS were consistent with a uniform distribution. Svensson et al. (2010) reached a similar conclusion using a larger sample of GOODS and PANS core-collapse SNe. For more local SN samples, Kelly et al. (2008) measured the fractional flux distributions for Type II and Ib/c SNe and found that Type Ic SNe are consistent with being drawn from the same fractional flux distribution as LGRBs, whereas Type II and Ib SNe uniformly track the light of their hosts. They suggest that LGRBs and Type Ic SNe share a common progenitor, and that a factor such as metallicity may determine whether core-collapse results in a LGRB with an associated Ic-BL SNe or a normal Type Ic SNe with no associated LGRB. However, recent work has found that LGRBs and Ic-BL SNe occur in host galaxies with high star formation rate density and that this preference cannot be due to metallicity (Kelly et al. 2014). Finally, the LGRB fractional flux distribution has also been compared to that for super-luminous supernovae yielding a result that is suggestive of agreement (Lunnan et al. 2015). With the growing samples of SNe, comparative studies of SN and LGRB environments have become limited by the small sample of LGRBs from the studies carried out a decade ago (Bloom et al. 2002; Fruchter et al. 2006). This situation can now be remedied with the much larger sample of *Swift* LGRBs.

Here we present the first uniform analysis of *HST* follow-up observations of the host galaxies of about 100 *Swift* LGRBs collected over the last decade. The goal of this analysis is twofold. First, to investigate the offset and fractional flux distributions for a much larger and more uniform sample of LGRBs than previously possible. Second, to provide a much larger comparison sample for future studies of the locations of

other astrophysical transients, such as various SN types. The structure of our paper is as follows. In Section 2 we present the details of the host galaxy and afterglow observations, data analysis, and astrometric matching. In Section 3 we discuss our offset, fractional flux, and host assignment methodologies, and present our measurements. In Section 4 we present our resulting offset and fractional flux distributions, and in Section 5 we discuss potential trends with redshift, the relationship between fractional flux and offset, and implications for LGRB progenitors. We conclude with a summary in Section 6.

In this paper we use $H_0 = 67 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.32$, and $\Omega_\Lambda = 0.68$ (Planck Collaboration et al. 2014), as well as AB magnitudes (Oke & Gunn 1983), corrected for Galactic extinction (Schlafly & Finkbeiner 2011).

2. OBSERVATIONS AND ASTROMETRY

2.1. *HST* Observations and Reduction

To locate the LGRBs within their host galaxies we utilize high resolution *HST* images. We primarily use Wide Field Camera 3 (WFC3) and Advanced Camera for Surveys (ACS) data, but also Wide Field Planetary Camera 2 (WFPC2) and Near Infrared Camera and Multi Object Spectrometer (NICMOS) data when the former are not available. Our sample is composed of 100 LGRBs observed over the last decade (August 2004 to July 2013) under multiple programs (see Table 1 for program ID numbers). We perform the first uniform analysis of these data and utilize them for the purpose of investigating the locations of LGRBs within their hosts and their relation to the underlying host light distribution. This sample has the significant advantage of largely consisting of bursts discovered by the *Swift* satellite and thus does not suffer from any potential biases that may be associated with combining data from multiple satellites as was done in previous studies. About 86% of the bursts in the sample have measured redshifts ($0.03 \lesssim z \lesssim 9.4$) and the *HST* imaging for these bursts span rest-frame UV and optical wavelengths (Figure 1). The wide redshift range allows us to investigate trends as a function of cosmic time. Unlike previous studies, the sample also includes a substantial number of dark LGRBs, which either lack optical afterglows or are much fainter than expected (GRBs 051022, 060719, 060923A, 061222A, 070306, 070521, 070802, 080207, 080325, 080607, 081109, 081221, 090404, 090407, 090417B, 090709A, 100413A, 100615A, and 110709B). Some are detected in the NIR while others are exclusively detected in the radio and X-ray bands.

We retrieved the *HST* observations from the MAST archive; for the ACS images we obtained charge transfer efficiency (CTE) corrected images, while for WFC3/UVIS data we used software from STScI to apply CTE corrections. We reduced the data using the *astrodrizzle* (Gonzaga et al. 2012) task in the STSDAS IRAF package utilizing recommended parameter settings for each instrument. By combining dithered exposures, this task enables the reconstruction of a higher resolution image than sampled by the instrumental point spread function. The task also applies distortion corrections to the images, which are critical for precise astrometric alignment. We use `final_pixfrac` = 0.8 and `final_scale` = 0.065", 0.02", 0.03", 0.05", and 0.15" per pixel for WFC3/IR, WFC3/UVIS, ACS, WFPC2, and NICMOS, respectively. In Table 1 we list for each LGRB the program ID, instrument, filter, and exposure time of the final drizzled image. In cases where there are multiple epochs of




Figure 1. Rest-frame wavelength versus redshift of the observations used to measure offsets and fractional flux for each LGRB. Only bursts with measured redshifts are included in this plot. Also shown are projected histograms of the redshift distribution and the distribution of rest-frame wavelengths probed by the *HST* observations.

imaging we list only the final epoch, unless an earlier epoch was chosen as the best host image. In Figure 2 we show the final drizzled image for each burst, with the location of the afterglow shown as determined by relative or absolute astrometry (Sections 2.3 and 2.4).

2.2. Afterglow Imaging

To precisely locate the LGRBs within their host galaxies requires the detection of an afterglow. About 85% of bursts in the sample have detected optical or NIR afterglows. We utilize the deepest, highest resolution afterglow images available to best match the quality of *HST* imaging. We primarily rely on publicly available afterglow images from the 8-m Very Large Telescope (VLT) and 8-m Gemini North and South telescopes. In 13 cases, an afterglow image is available from *HST*. We also use images from the *Swift* UV/Optical Telescope (UVOT), 6.5-m Magellan telescopes, Palomar 60-inch Telescope, Swope 40-inch Telescope, and the 8.2-m Subaru Telescope.


We use standard IRAF packages to analyze the afterglow data. We use the task `uvotimsum` in the HEASoft package to co-add *Swift*/UVOT images, combining multiple filters if necessary to obtain a high signal-to-noise ratio (S/N) afterglow detection. For *HST* afterglow images we use `astrodrizzle` as described in Section 2.1. For GRB

070306 we use the `eclipse` pipeline to reduce the NIR data from VLT/ISAAC. For GRBs 080325 and 090709A we use the MCSRED IRAF package to reduce the NIR data from Subaru/MOIRCS. For the remaining bursts without optical or NIR afterglows, we rely on radio or X-ray detections. We obtain the reported radio positions from the GCN circular archive. When an X-ray afterglow image is available from *Chandra*, we retrieve the image from the *Chandra* Data Archive. For each burst, we list in Table 1 the telescope and filter utilized to obtain the afterglow image that was used for astrometry.

2.3. Relative Astrometry


To precisely locate the afterglows on the *HST* images we use relative astrometry. We first identify common point-like sources between the afterglow and *HST* images, using `SExtractor` (Bertin & Arnouts 1996) to measure the source positions. We then match the coordinates with the IRAF task `ccmap`, using a second or third order polynomial to compute the astrometric solution. Using the afterglow images from the telescopes listed in Section 2.2 we generally obtain the following astrometric tie uncertainties, σ_{tie} (1σ):

- 6 mas using an *HST* detection of an optical or NIR afterglow




(a) Panel One

Figure 2. *Hubble Space Telescope* drizzled images of 100 LGRB host galaxies studied in this work. The circles (3σ) indicate the location of each GRB. North is up and East is to the left. Red, blue, green, cyan, and yellow circles indicate that afterglow positions were determined from optical/NIR, *Swift*/UVOT, *Chandra*, radio, and XRT images, respectively. Physical scales are given when the LGRB redshift is known.




(b) Panel Two

Figure 2. Continued




(c) Panel Three

Figure 2. Continued



(d) Panel Four

Figure 2. Continued



(e) Panel Five

Figure 2. Continued

- 30 mas using an optical or NIR afterglow detection from ground-based telescopes
- 100 mas using a UV or optical afterglow detection from *Swift*/UVOT
- 300 mas using an X-ray afterglow detection from *Chandra*

In a few cases, the lack of sufficient common sources for a robust astrometric tie necessitates the use of an intermediate image to locate the afterglow position on the *HST* image. This is the case for bursts detected with *Chandra*. We use the CIAO task `wavdetect` to measure the positions of X-ray sources. In these cases, the uncertainty associated with the match between the *Chandra* and intermediate ground-based images dominates the total astrometric tie uncertainty. Table 2 lists the number of common sources and astrometric tie uncertainty for each burst. Bursts with no listed astrometric tie uncertainty are those for which the best afterglow position is from XRT or for which we were not able to obtain an afterglow image. GRB 060418 is a unique case where the afterglow and host are well-detected in a single *HST* image and are sufficiently offset from each other such that their positions are not biased by light contribution from each other (later epoch imaging revealed no host emission at the afterglow position). Therefore astrometric matching is not necessary for this burst.

2.4. Absolute Astrometry

In cases where the afterglow is detected only in the radio band, or when the X-ray observations do not contain any common sources for relative astrometry, absolute astrometry is performed by matching the late-time *HST* images to the 2MASS astrometric system. This is mainly the case for dark GRBs. For the two GRBs where a radio position was used (090404 and 100413A), the uncertainty in the afterglow position with respect to the *HST* image is dominated by the astrometric accuracy of those radio positions, $\sim 0.5''$. Similarly, when using absolute astrometry to locate the X-ray afterglow of GRB 090407, the error is dominated by the astrometric accuracy of *Chandra*, $\sim 0.4''$.

3. MEASUREMENT METHODOLOGY

3.1. Offset Measurements

Once the afterglow and *HST* images are registered, we identify the host galaxy in the *HST* image. First we measure the afterglow centroid using `SExtractor` to determine the afterglow position relative to the *HST* image. The afterglow positional uncertainty (σ_{AG}) is estimated as $\frac{FWHM}{2(S/N)}$. In some cases, it is apparent in the afterglow image that there is an underlying extended object, presumably the host galaxy. To reliably measure the centroid of the afterglow we perform image subtraction, using the ISIS software package (Alard 2000), relative to a late-time image of the host ideally from the same telescope and in the same filter. Once the afterglow position is determined, we use a uniform host assignment procedure discussed in Section 3.3. Here we continue our discussion of the offset measurement procedure assuming a host galaxy has been identified.

We measure the host galaxy center and corresponding uncertainty using `SExtractor`. We use a uniform definition of the host center as the brightest region of the galaxy (or peak

of the light distribution) determined by sequentially increasing the threshold for detection in `SExtractor`. The statistical uncertainty on the centroid determined by `SExtractor` does not take into account the larger systematic uncertainties caused by the fact that LGRB hosts tend to have irregular morphologies. Instead, the uncertainty on the host center (σ_{host}) is estimated by taking the quadrature sum of the standard deviation in both directions of the positions measured by multiple runs of `SExtractor` using varying thresholds for detection from 3σ to the highest threshold that yields a detection.

Using the afterglow position and host center we calculate the projected angular offset (R) between the afterglow position and host center. The total uncertainty on the angular offset (σ_R) is taken to be the quadrature sum of σ_{tie} , σ_{AG} , and σ_{host} . We also calculate the projected physical offsets using the measured redshift for each burst. For bursts with no measured redshift we use the fact that at $z \gtrsim 0.5$, typical of most LGRBs, the angular diameter distance is approximately constant so that we can use a conversion factor of about $8 \text{ kpc}''$. The projected angular and physical offsets and corresponding uncertainties are listed in Table 2. Bursts for which no offsets are listed indicate either bursts with well-localized afterglows but no detected host (see Section 3.3 for a discussion of distinguishing between no host detection and a large offset) or bursts for which we only have an XRT position or were unable to access the afterglow image. The uncertainties on the XRT positions are typically $\gtrsim 1''$ in radius, too large to enable a reliable host association.

3.2. Host Galaxy Sizes, Normalized Offsets, and Magnitudes

As our interest lies in the positions of LGRBs relative to the light of their host galaxies, we normalize the offsets by the sizes of the hosts. Here we use the half-light radius (R_h) as measured by `SExtractor` (with `FLUX_RADIUS` = 0.5). We also measure the 80% light radius (R_{80}) for comparison with previous studies of LGRBs and SNe. In Table 2 we list the half-light radii and host-normalized offsets (R_{norm}) for each burst.

We also measure the apparent magnitudes of the host galaxies, and corresponding uncertainties, using the `MAG_AUTO` estimate from `SExtractor` which uses Kron apertures. We obtain the magnitude zeropoints from STScI where tabulated, or from the photometry keywords in the *HST* image headers. The magnitudes corrected for Galactic extinction (Schlafly & Finkbeiner 2011) and corresponding uncertainties are listed in Table 2 for each burst with an associated host galaxy.

3.3. Host Galaxy Assignment

To assign host galaxies to each LGRB we calculate the probability of chance coincidence, for several candidate extended objects surrounding the afterglow position, given the observed number density of field galaxies from deep optical and NIR surveys (Hogg et al. 1997; Beckwith et al. 2006; Metcalfe et al. 2006). Nominally we assign the candidate with the lowest probability of chance coincidence as the host. For candidates at small offsets from the afterglow position, such that the afterglow position is essentially coincident with the candidate, this candidate will in most cases have the lowest probability of chance coincidence. Bursts for which there is no coincident candidate require further investigation to distinguish between the scenario where the host was not detected and the scenario where the burst occurred at a large offset from a neighboring galaxy. We discuss these cases in detail

in Section 3.3.1. Following the methodology of Bloom et al. (2002) we determine the probability of chance coincidence (P_{cc}) using the following equation:

$$P_{cc} = 1 - e^{-\pi R_e^2 \sigma(\leq m)} \quad (1)$$

where $\sigma(\leq m)$ is the observed number density of galaxies brighter than magnitude m , and R_e is the effective radius. For optical observations we use the galaxy number densities given by Equation 1 of Berger (2010), whereas for NIR observations we use the H -band number counts measured by Metcalfe et al. (2006). As discussed by Bloom et al. (2002), the appropriate value of R_e depends on the offset, size of the galaxy, and uncertainty on the afterglow position. Following Bloom et al. (2002) we take R_e to be the maximum of $(3\sqrt{\sigma_{tie}^2 + \sigma_{OT}^2}, \sqrt{R^2 + 4R_h^2})$. In other words, if the localization is precise, such that the first term is much smaller than the offset and galaxy size, then the second term dominates. On the other hand, if the GRB localization is poor the positional uncertainty dominates over the offset and galaxy size, and its radius becomes the appropriate value of the effective radius for determining P_{cc} . In the majority of cases ($\sim 85\%$) the localizations are sufficiently precise such that the second term dominates over the first. As discussed in Section 2.3, σ_{tie} is typically $\gtrsim 100$ mas for bursts where we use afterglow detections from *Swift*/UVOT or *Chandra*. In these cases, the GRB positional uncertainty can dominate over the offset and galaxy size.

We list the values of P_{cc} in Table 2. For the bursts with coincident extended objects, P_{cc} tends to fall in the range 10^{-3} to 5×10^{-2} . Therefore we do not expect significant contamination of our host galaxy sample by unrelated galaxies along the line of sight.

3.3.1. Bursts With No Coincident Host Candidate

The subset of bursts that lack a coincident extended object must be examined further to determine if the object with the lowest P_{cc} can realistically be considered the host galaxy. Relying simply on P_{cc} may result in the inclusion of unrealistically luminous objects in the sample. We identify as possible host candidates the nearest extended objects with $P_{cc} \lesssim 0.1$. Given the measured redshift of the burst, we can determine whether the luminosity of the host candidate is consistent with the range of LGRB host luminosities observed at that redshift. It may be more likely that the true host lies beyond the magnitude limit of our observations and was not detected. However, it is important to avoid biasing our results by eliminating potential large offsets. Here we discuss our assessment of bursts whose host candidates satisfy $P_{cc} \lesssim 0.1$ but are offset by significantly more than the half-light radius of the host candidate.

To carry out the assessment of non-coincident host candidates with $P_{cc} \lesssim 0.1$ we compare their luminosities to the luminosities of the coincident host galaxies. We convert our measured apparent magnitudes to rest-frame luminosities in units of L^* . Given the redshift, we determine the absolute magnitude and using measured values of L^* from galaxy surveys (Faber et al. 2007; Marchesini et al. 2007; Bouwens et al. 2015) we calculate L/L^* for each host galaxy. For bursts with $z \lesssim 1.7$ we use the measured value of L^* in the B -band for the blue galaxy sample analyzed by Faber et al. (2007). In the range $1.7 \lesssim z \lesssim 3.2$ we use L^* values in the B -band from Marchesini et al. (2007) using their $U-V < 0.25$ mag galaxy subsample. For bursts with $z \gtrsim 3.2$ we use L^* corresponding

to UV bands as measured by surveys of Lyman-break galaxies (Bouwens et al. 2015). These were chosen so that the wavelength regime in which the L^* values are appropriate correspond roughly to the rest-frame wavelengths probed by our observations in these redshift bins. We do not perform a K -correction to a common band.

We plot the luminosities as a function of redshift for the full sample, including coincident and non-coincident candidates, in the left panel of Figure 3. The right panel shows the luminosities as a function of host-normalized offset. The luminosities of the robust host galaxy associations span $\sim (0.01 - 2)L^*$, whereas some host candidates at relatively large offset and high P_{cc} have luminosities ranging from $\sim (3 - 100)L^*$, inconsistent with the distribution for bursts with coincident host galaxies and indicating that they are unrelated galaxies. The luminosity of the host candidate for GRB 060526 is $\sim 100L^*$ assuming it is at the redshift of the GRB, much larger than expected for a GRB host. We therefore reject this association. We also reject the candidates for GRBs 060927, 070125, 071031, and 091208B because their luminosities are unreasonably high given the observed distribution for securely assigned hosts. In the case of GRB 080928, there are two candidates with $P_{cc} \lesssim 0.1$ where the object with $P_{cc} \approx 0.023$ has a luminosity of $\sim 4L^*$ and the object with $P_{cc} \approx 0.035$ has a luminosity of $\sim L^*$. We therefore accept the lower luminosity galaxy as the host. We also accept as the host the candidate for GRB 081008 due to its reasonable luminosity of $\sim 0.6L^*$. The left panel of Figure 3 also shows upper limits on the luminosities of the hosts where we are confident the host was not detected due to the lack of any nearby host candidates with $P_{cc} \lesssim 0.1$. Upper limits for the cases with rejected host candidates discussed above are also shown.

Furthermore, we note that there is no trend of host luminosity versus host-normalized offset. There is no reason to suspect from predictions that bursts at large or small offsets occur only in bright or faint galaxies. As shown in Figure 3, the inclusion of the rejected host candidates creates an artificial trend because the only galaxies at higher normalized offset that have $P_{cc} \lesssim 0.1$ are bright galaxies. Therefore we are confident these rejected candidates are unrelated galaxies and the true host was not detected. Indeed, the resulting upper limits on the luminosities of these hosts are not surprising given the upper limits we calculate for unambiguous non-detected hosts. Upon completing our host assignment procedure we obtain a sample size of 71 LGRBs for which we make offset and fractional flux measurements.

3.4. Fractional Flux Measurements

Studying the locations of LGRBs within their hosts can be expanded beyond an analysis of the offset distribution. In many cases where the host's morphology is irregular, the location of the host center is unclear and the offset from an empirically defined center may not be meaningful. Furthermore, the host galaxy size defined by the half-light radius does not capture the complex morphology of many of the hosts. While measuring offsets is useful for comparing the locations of LGRBs with predicted profiles of star formation, we would like to go one step further and ask whether or not LGRBs are spatially coincident with sites of massive star formation. This is possible using a method, less sensitive to galaxy morphology than offsets, of comparing the brightness at the LGRB location relative to the entire host light distribution (Fruchter et al. 2006). We calculate the fraction of the total flux from the host galaxy that is contained in pixels fainter than or equal to




Figure 3. *Left:* Host galaxy rest-frame luminosity in units of L^* as a function of redshift for assigned host galaxies (red), rejected host candidates (green), and non-detected hosts (blue). The data point area is proportional to P_{cc} where a smaller point means a smaller P_{cc} . The red dotted line indicates a luminosity of L^* , and the dotted cyan line connects the two possible host candidates for GRB 080928 (see Section 3.3.1). We do not include bursts with unknown redshift. *Right:* Host galaxy luminosity versus host-normalized offset. As expected this is a scatter plot for the robust host associations. The correlated tail caused by the rejected candidates is an indication that they are unrelated galaxies.

the flux at the LGRB location. The resulting "fractional flux" is a statistic that measures the brightness of the burst site compared to the entire galaxy. A fractional flux value of 1 means that the burst occurred on the brightest pixel in the galaxy.

We measure the fractional flux for each LGRB using the following procedure. First we determine the flux value at the LGRB position. When the 1σ error circle defined by the uncertainty on the LGRB position as determined by the quadrature sum of σ_{tie} and σ_{AG} spans less than 1 pixel, we simply take the flux value of that pixel. When the error circle encompasses more than one pixel but is smaller than the PSF, we measure an average flux within the 1σ error circle weighted by the fractional area encompassed by a given pixel. These scenarios are relevant for most bursts where we use afterglows from ground-based optical/NIR telescopes and *HST* (see Section 2). Using *SExtractor* we extract the host galaxy pixels using a threshold of 1 σ above the sky background and then calculate the fraction of the total flux in pixels fainter than or equal to the average flux of the LGRB error region (the fractional flux).

When the error circle is larger than the PSF, the galaxy brightness as a function of position could vary significantly so that the uncertainty in the position of the LGRB may have a large impact on the fractional flux. This scenario is often encountered when using optical afterglow detections from *Swift*/UVOT, X-ray afterglows from *Chandra*, or radio afterglow detections, where the error regions can encompass tens to hundreds of pixels. In some cases, these pixels include both the faintest and brightest pixels of the host galaxy. In this regime, it is unclear whether the fractional flux determined

as described above remains a meaningful quantity. Here we show that for bursts with error circles larger than the PSF, a different weighting is required to properly assess the fractional flux in these cases.

To assess this, we employ a different procedure for determining the fractional flux, taking into account the Gaussian error distribution associated with each burst. As before we extract the galaxy pixels using *SExtractor*. Each pixel in the image has an associated probability for hosting the burst, governed by the 2D Gaussian distribution that takes into account the astrometric tie and afterglow positional uncertainties. For all pixels encompassed by the error circle, a corresponding fractional flux value can be calculated. Pixels not associated with the galaxy will have fractional flux values of zero. The posterior probability distribution for the fractional flux can then be constructed by weighting each value by its corresponding Gaussian probability. The fractional flux and corresponding uncertainty are estimated as the mean and standard deviation of this distribution.

By comparing the fractional flux values for bursts with large error regions with the well-localized bursts, we find that there exists a considerable bias towards lower fractional flux values in the former. We hypothesize that not only is the absolute size of the error circle important but also the error circle size relative to the galaxy size. If the galaxy is much larger than the error circle, even a large error circle in an absolute sense can yield a well-constrained fractional flux value. If a relatively well-localized burst occurs in a small galaxy, the fractional flux value may not be well-constrained. Figure 4 illustrates this bias by plotting fractional flux versus the ratio




Figure 4. *Left:* Fractional flux versus the ratio of error circle area to galaxy area. Green points with error bars represent bursts with error circles larger than the *HST* PSF. *Right:* The fractional flux distribution binned into two bins separated at an area ratio of 0.1. In both plots it is apparent that there is a bias to lower fractional flux values when the error circle covers a substantial fraction of the galaxy area. In the analysis in Section 4.2 we only use bursts with a ratio of $\lesssim 0.1$

of error circle area to galaxy area (left panel). The right panel of Figure 4 shows the cumulative distributions for two subsets of the data divided at an area ratio of 0.1. Of the bursts with area ratios $\gtrsim 0.1$, there are none with fractional flux values greater than ~ 0.6 , while bursts with an area ratio of $\lesssim 0.1$ span the full range of fractional flux values. This bias can be understood by noting that as the error region increases relative to the galaxy size, there is greater probability that the burst occurred on background sky pixels. In general we find that for error regions large in an absolute or relative sense there is a bias towards lower fractional flux values. To avoid biasing our results we remove from consideration the bursts with area ratios of $\gtrsim 0.1$, resulting in a sample size of 50 LGRBs. By removing these bursts we are not concerned about selecting against bursts with intrinsically low fractional flux values. Here we are removing a bias associated with the measurement of fractional flux when burst localization is poor.

In Table 3 we list the fractional flux values and area ratios for each LGRB. Bursts with no fractional flux value are either bursts for which a host was not detected or a sufficient afterglow position was not available.

4. RESULTS

We have compiled the largest sample of LGRB offsets and fractional flux measurements presented to date. Our aim is to use them to understand where within their host galaxies LGRBs occur and what the results from a large sample indicate about the progenitors. Here we present the results of our offset and fractional flux distribution measurements summarized in Tables 2 and 3.

4.1. Offset Distribution

In this section we present the measured physical and host-normalized offset distributions. We also present the results of

an analysis of the offset uncertainties to understand how they affect the offset distributions. Each offset measurement has an associated uncertainty (σ_R) dependent on σ_{tie} , σ_{AG} , and σ_{host} (Section 3.1). Due to the inherent non-Gaussian nature of offsets and the non-uniform uncertainties, we employ a Monte Carlo approach to assess the uncertainty on the resulting distributions of offsets. At each iteration a random offset (x) was drawn for each burst from its offset probability distribution defined by the measured offset (R) and uncertainty (σ_R). For bursts with $R/\sigma_R \lesssim 5$ we use a Rice distribution to represent the probability distributions, defined by:

$$p(x|R, \sigma_R) = \frac{x}{\sigma_R^2} \exp \left[-\frac{(x^2 + R^2)}{2\sigma_R^2} \right] I_0 \left(\frac{xR}{\sigma_R^2} \right) \quad (2)$$

where I_0 is the zeroth order modified Bessel function of the first kind. For $R/\sigma_R \gtrsim 5$, the Rice distribution can be approximated as a Gaussian. We make this approximation for numerical ease. In all cases discussed in the following sections we use 10,000 iterations.

4.1.1. Physical Offset Distribution

In Figure 5 we show the cumulative distribution of projected physical offsets, calculated as described in Section 3.1. The final sample size included in this distribution is 71 LGRBs, the subset of the original sample where offset measurement is possible. The distribution ranges from offsets of 0.075 to 14.4 kpc with the median at 1.42 kpc. About 90% of bursts occur within ~ 5 kpc. The distribution is remarkably smooth and has bursts representing the full range of offsets. There are few gaps where we do not find bursts, with one notable gap at $\sim 7 - 14$ kpc. For comparison we also show the distribution of physical offsets for the small sample of 20 LGRBs measured by Bloom et al. (2002). The two distributions agree exceptionally well. A Kolmogorov-Smirnov (KS)




Figure 5. *Left:* Cumulative distributions of the projected physical offsets of 71 LGRBs from this work (red) and Bloom et al. (2002) (B02, blue). We also show the distributions for Type Ib/c (cyan) and Type II (green) SNe from Prieto et al. (2008). The shaded region displays the results of our Monte Carlo assessment of the uncertainties on the offsets in the form of a 2D histogram. *Right:* Similar to the left panel, except here we plot the sample of 58 bursts with $\sigma_{R_{\text{phys}}} \lesssim 1.0$ kpc to reduce any bias caused by bursts with a large offset uncertainty (See Figure 6). The results are essentially unchanged.

test yields a p-value of 0.94. We also show in Figure 5 the results of our Monte Carlo simulation (taking into account the offset uncertainties) by plotting a 2D histogram showing the density of points from the resulting cumulative distributions generated at each iteration. Darker regions indicate a higher density of points, in other words a high fraction of synthetic distributions pass through that region. The apparent shift of the distribution from the measured distribution towards higher offset values is indicative of the skew to higher offsets resulting from the fact that an offset is a positive-definite quantity described by a Rice distribution. The shift is dominated by the ~ 10 LGRBs for which the uncertainty is larger than the offset.

It is important to carefully consider the effects of the uncertainties on the distribution of offsets because offsets with uncertainties large relative to the offset itself have probability distributions skewed to larger offset. Given a large uncertainty, the measured offset is more likely to be large due to the large area encompassed by the error circle at large offset. As $\sigma_{R_{\text{phys}}}$ should not be correlated with the intrinsic offset, if the true offset distribution extended to small offsets, this effect would cause a bias against finding them when the uncertainties are large. In Figure 6 we plot $\sigma_{R_{\text{phys}}}$ versus R_{phys} . Of the 13 bursts in the sample with $\sigma_{R_{\text{phys}}} \gtrsim 1.0$ kpc, all are at measured offsets $\gtrsim 0.5$ kpc, whereas the remaining bursts with $\sigma_{R_{\text{phys}}} \lesssim 1.0$ kpc span the full range of offsets from 0.075 to 14 kpc. Since there is no relationship between the true offset and $\sigma_{R_{\text{phys}}}$, the range of offsets found with large uncertainty should not be significantly different from the range found with small uncertainty. The fact that we only measure large offsets when we have large uncertainty is indicative of the above bias. We note that even at relatively small uncertainty we may still be missing bursts at very small offsets because any amount of uncertainty prevents the measurement of arbitrarily small offsets.

To reduce this bias, we make a quality cut on our physical offset sample by also plotting in Figure 5 the distribution for bursts with $\sigma_{R_{\text{phys}}} \lesssim 1.0$ kpc, resulting in a sample of 58 LGRBs. The resulting distribution has a median at 1.27 kpc. The mean of the distribution of medians for each iteration of the Monte Carlo simulation is 1.34 kpc with a 90% confidence interval of 1.17 – 1.51 kpc.

We also compare the distribution of projected physical offsets to the distributions for Type II and Ib/c SNe. KS tests with the supernovae samples of Prieto et al. (2008) yield p-values of 1.33×10^{-6} and 1.53×10^{-7} for the Type Ib/c and II SNe, respectively. LGRBs occur significantly closer to the centers of their host galaxies in a physical sense than SNe. To disentangle the effects of galaxy size differences between LGRB hosts and SNe hosts, we revisit supernova comparisons in the next section where we analyze the LGRB host-normalized offsets.

4.1.2. Host-Normalized Offset Distribution

Normalizing the offsets by the host galaxy sizes enables a fair comparison across the LGRB sample and with the SNe samples. In Figure 7 we plot the cumulative distribution of host-normalized offsets (R/R_h). The distribution ranges from about 0.04 to 7.0. The median of the distribution is 0.79 and about 90% of the bursts occur within a host-normalized offset of ≈ 2.2 . The distribution is overall reminiscent of an exponential disk profile, the expected surface brightness profile of star forming disk galaxies. For comparison we also plot the predicted distribution of host-normalized offsets if LGRB locations exactly trace an exponential disk profile. Although the shapes are overall consistent between our measured distribution and that of an exponential disk, there is a notable shift to lower offsets seen in our distribution. Again, to compare with the previous offset sample of LGRBs we show the host-normalized offsets from Bloom et al. (2002). A KS test between this sample and our sample gives a p-value of 0.50.




Figure 6. *Left:* Uncertainty in physical offset ($\sigma_{R_{\text{phys}}}$) versus physical offset (R_{phys}). The green line marks $\sigma_{R_{\text{phys}}} = R_{\text{phys}}$. *Right:* Uncertainty in host-normalized offset ($\sigma_{R_{\text{norm}}}$) versus host-normalized offset (R_{norm}). Similarly, the green line marks $\sigma_{R_{\text{norm}}} = R_{\text{norm}}$. Since $\sigma_{R_{\text{norm}}}$ and $\sigma_{R_{\text{phys}}}$ are not related to the intrinsic offset, the apparent lack of small offsets with large uncertainty (the gap in the upper left quadrant of both plots) is indicative of a bias to large offset associated with bursts that have poor localizations. The dashed blue lines at $\sigma_{R_{\text{phys}}} = 1$ kpc and $\sigma_{R_{\text{norm}}} = 0.5$ mark the locations of our cuts.

As a result of the small sample size, Bloom et al. (2002) concluded that the LGRB host-normalized offsets were consistent with being drawn from the exponential disk profile. With our larger sample we can rule this out at a significance level of 0.01. Given the irregular morphologies of LGRB host galaxies this result is not surprising even if LGRBs trace star formation.

As with the physical offsets, we also show in Figure 7 the uncertainty region determined from our Monte Carlo procedure. Again, we carefully consider the effects of the uncertainties on the distribution of host-normalized offsets to avoid the aforementioned bias associated with the offset measurement of bursts with large offset uncertainties. In Figure 6 we also plot $\sigma_{R_{\text{norm}}}$ versus R_{norm} . Of the 20 bursts in the sample with $\sigma_{R_{\text{norm}}} \gtrsim 0.5$, all are at measured offsets $\gtrsim 0.4$, whereas the remaining bursts with $\sigma_{R_{\text{norm}}} \lesssim 0.5$ span the full range of offsets from 0.04 to 7.0. As with the physical offsets, our inability to measure small offsets when we have large uncertainty is indicative of a bias. Again, it is likely the case that even at relatively small uncertainty we may still be missing bursts at very small offsets.

As before, we make a quality cut on our host-normalized offset sample by restricting the sample to bursts with $\sigma_{R_{\text{norm}}} \lesssim 0.5$, resulting in a sample of 52 LGRBs. The resulting distribution has a median at 0.54. In Figure 8 we show the individual probability distributions of each LGRB used in the Monte Carlo simulation. The mean of the distribution of medians for each iteration of the Monte Carlo simulation is 0.67 with a 90% confidence interval of 0.58 – 0.76. We plot the distribution and Monte Carlo results in Figure 9. A KS test with the exponential disk model yields a p-value of 2.6×10^{-4} . Furthermore, only $\sim 4\%$ of the Monte Carlo synthetic distributions have p-values > 0.05 when compared to an exponen-

tial disk profile, meaning that we can rule out the hypothesis that LGRB host-normalized offsets are drawn from an exponential disk. As another way to view these results we also show in Figure 8 the sum of the individual probability distributions of each LGRB. The summed distribution effectively produces a smoothed histogram of the offsets and shows a clear shift to smaller offsets from the probability distribution for an exponential disk profile. LGRBs are apparently highly concentrated in the inner parts of their hosts. In particular, they are more concentrated than the radial light distributions of their hosts; namely, 50% of LGRBs occur within 67% of a half-light radius. Stated differently, about half of all LGRBs occur within a region that contains only 33% of the underlying distribution of light in their host galaxies. The possible bias caused by the inability to find bursts at very small offset mentioned above would act in the opposite direction of the striking result found here, indicating that the distribution may be even more concentrated than we find here.

A striking feature of the distribution in Figure 9 is the apparent tail to large offset extending from a gap at $R_{\text{norm}} \sim 1.6 - 3.3$. While it is not unexpected, given the rough similarity of our distribution to an exponential disk, to find bursts at large offset, it is somewhat surprising that $\sim 10\%$ of LGRBs appear to be located at $R_{\text{norm}} \gtrsim 3$. These bursts include GRBs 050820, 081008, 080928, 060418, and 060505. We revisit the properties of these bursts and their assigned hosts. The host of GRB 050820 has a morphology consisting of a bright core with a diffuse tail with the GRB located on this diffuse emission, leading to a host-normalized offset of 3.8 from the bright core. The burst’s location on the underlying galaxy emission and the P_{cc} of 0.01 do not give reason to doubt the host association, the large offset being due to the unusual morphology. As discussed in Section 3.3.1 we assigned as the host




Figure 7. Cumulative distributions of LGRB host-normalized offsets for our sample (red) and the sample of Bloom et al. (2002) (B02, blue) shown on linear (left) and log (right) x -axis scales. Also shown are the distributions for Type Ib/c (cyan) and Type II (green) SNe from Kelly & Kirshner (2012) and SLSNe (orange) from Lunnan et al. (2015). We also plot the distribution expected for an exponential disk profile (magenta). The shaded regions depict the uncertainties on our offset measurements determined using a Monte Carlo simulation. We find an offset distribution qualitatively similar to an exponential disk profile but statistically inconsistent.




Figure 8. *Left:* Probability distributions for each LGRB with $\sigma_{R_{\text{norm}}} \lesssim 0.5$ given its measured offset and offset uncertainty calculated using Equation 2. Narrow peaks indicate well-localized bursts while broader distributions indicate poorer localizations. *Right:* Sum (red) of the individual probability distributions (left) effectively producing a smoothed histogram of the offsets. In magenta we show the probability distribution for an exponential disk. The summed distribution is clearly offset to lower offsets than what would be expected for a population tracing an exponential disk.

of GRB 080928 a non-coincident galaxy with a luminosity consistent with the observed distribution of LGRB host luminosities. However, we cannot rule out the possibility that the host was not detected and that the assigned candidate is in-

stead the source of an intervening absorption system observed in the afterglow (Vreeswijk et al. 2008), casting doubt on the large inferred offset. GRB 081008, another burst with a host assigned based on our luminosity analysis, has a large normal-




Figure 9. Similar to Figure 7 except here we show the distribution of host-normalized offsets only for bursts with $\sigma_{R_{\text{norm}}} \lesssim 0.5$. The distribution is notably shifted to lower offset with a median of 0.54. This is because we have removed bursts with large offset uncertainties which bias the distribution to larger offsets (See Figure 6).

ized offset of ≈ 7 and $P_{\text{cc}} \approx 0.07$. While the host assignment matches our criteria, we acknowledge the possibility that the assignment is not correct. For GRB 060418 we assign as the host the brighter of the two nearby host components identified by Pollack et al. (2009). Although these components are distinct from each other (see Figure 2), the assignment of one or the other has little effect on the offset and no effect on the fractional flux. GRB 060505 is a controversial $z = 0.089$ burst that did not have an associated supernova as expected for a low- z LGRB leading some to argue in favor of a short GRB scenario (Ofek et al. 2007). However, given its location in a bright, low-metallicity HII region of its host, others argue in favor of 060505 as a LGRB (Thöne et al. 2014). In this case, 060505 is an example of a LGRB occurring at large normalized offset (3.6) and at high fractional flux (0.99) in the outskirts of its host. It is interesting to speculate that the other bursts at large offset may be higher redshift versions of 060505 where underlying HII regions are too faint to be detected. For the reasons discussed here, the large offset tail in the distribution of host-normalized offsets may not be real. However, it does not affect the overall conclusion that LGRBs as a population are more centrally concentrated than the underlying light distributions of their hosts.

We compare our distribution of LGRB host-normalized offsets with those measured for supernovae by Kelly & Kirshner (2012), also shown in Figure 9. KS tests with the Type Ib/c and Type II SNe yield p-values of 1.3×10^{-2} and 5.2×10^{-5} , respectively. While the measured distribution for Type Ib/c and our sample of LGRBs are statistically inconsistent, about 80% of Monte Carlo synthetic distributions have p-values > 0.05 when compared to Type Ib/c SNe. This percentage is $\sim 1\%$ when compared to Type II SNe. When taking into account the uncertainties, the distributions for LGRBs and Type Ib/c SNe are therefore consistent. They both are much more centrally concentrated than Type II SNe and the exponential disk profile. Comparing our LGRBs to the distribution mea-

sured for super-luminous supernovae (SLSNe; Lunnan et al. 2015) we find a KS test p-value of 0.17. This comparison, though, is limited by the small SLSNe sample size.

4.1.3. Offset Distribution Summary

We find that our sample of LGRBs spans the range 0.075 to 14.4 kpc (mean of Monte Carlo distribution of medians = 1.34 kpc) in physical offset from the centers of their host galaxies. When appropriately accounting for uncertainties and removing bursts with $\sigma_{R_{\text{norm}}} \gtrsim 0.5$, we find that LGRB host-normalized offsets are considerably smaller and more centrally concentrated (mean of Monte Carlo distribution of medians = 0.67, with 90% confidence interval of 0.58 - 0.76) than what would be expected if LGRBs traced an exponential disk profile. We can rule out at high significance that LGRB offsets are drawn from an exponential disk distribution and their distribution is inconsistent with Type II SNe. Accounting for uncertainties, the LGRB offset distribution we find is consistent with that for Type Ib/c SNe. LGRBs prefer the central locations in their hosts, being more concentrated than the underlying host light distributions. Table 4 provides a summary of the KS p-values we find for comparisons between our sample of LGRB host-normalized offsets and other types of stellar explosions and in Table 5 we show the medians.

4.2. Host Light Distribution

In Figure 10 we show the cumulative distribution of fractional flux for the 49 bursts in our sample with error circle to galaxy area ratios of $\lesssim 0.1$ (left panel). Ideally, we would like to assess whether or not LGRBs are spatially coincident with bright rest-frame UV regions of their hosts, which trace young, massive stars. As shown in Figure 1 our observations primarily probe the rest-frame optical emission, which may probe older stellar populations. Here we compare the fractional flux distribution for the $\sim 20\%$ of the sample that probes rest-frame UV to the distribution that probes

rest-frame optical. The right panel of Figure 10 shows the resulting distributions when dividing the sample by wavelength into two bins separated at $\lambda_{\text{rest}} = 5000 \text{ \AA}$. The rest-frame optical and blue/UV distributions agree well, which we interpret as indicating that there is no bias in using rest-frame optical observations as a proxy for rest-frame UV. If there was a separate older stellar population, we would not expect to find LGRB positions correlated so strongly with both the optical and UV light. In other words, the optical and UV light are probing the same population of stars. We can then interpret our fractional flux measurements as reflecting the relationship between location and the host UV emission.

As shown in Figure 10 the fractional flux distribution for our sample spans the full range from 0 to 1 with 90% of values falling in the range 0.03 to 0.98. The median is 0.75 and nearly 80% of bursts have fractional flux values greater than 0.5. This strong preference towards higher fractional flux values indicates that LGRBs are typically located on some of the brightest regions of their hosts, and lends further support to the idea that LGRBs originate from very massive stars. This is in broad agreement with previous claims (Fruchter et al. 2006; Svensson et al. 2010). We also show in Figure 10 the distribution of fractional flux for the sample of 30 LGRBs from Fruchter et al. (2006). A KS test between the two samples yields a p-value of 0.09. This indicates that while our sample and the Fruchter et al. (2006) sample are marginally inconsistent, we cannot rule out that they are drawn from the same distribution. Nevertheless, the sample of Fruchter et al. (2006) has a higher median and general inclination towards higher fractional flux values. This slight discrepancy is probably not caused by effects intrinsic to the LGRB population but rather by systematic differences between the two samples. We also show the Fruchter et al. (2006) sample with the wavelength binned distributions in Figure 10. The slight shift of the rest-frame blue/UV distribution towards the distribution of Fruchter et al. (2006), mostly comprised of rest-frame UV observations, may indicate the discrepancy is a bandpass effect. We address another potential effect in Section 5.

We also compare our measured distribution to the corresponding distributions measured for CCSNe (Svensson et al. 2010), SLSNe (Lunnan et al. 2015), and Type Ic and II SNe (Kelly et al. 2008). KS tests between our sample, the SLSNe sample, the Type Ic SNe sample, the Type II SNe sample, and the CCSNe sample yield p-values of 0.52, 0.79, 5.4×10^{-5} , and 0.03, respectively. We cannot rule out that our sample of LGRBs and the SLSNe sample are drawn from the same distribution. Interestingly, our sample overlaps the SLSNe distribution at fractional flux values of $\gtrsim 0.8$. The very good agreement between the distributions for our sample and the Ic SNe sample is consistent with the fact that Ic SNe, particularly Ic-BL, are the only types of SNe observed to be associated with LGRBs. The distributions of fractional flux for the CCSNe, mostly consisting of Type II SNe, and the Type II SNe sample are clearly distinct from our LGRB distribution, strong evidence that Type II SNe and LGRBs have different progenitors. We summarize the KS p-values for our comparisons in Table 4.

4.3. Galaxy Sizes

As it pertains to our knowledge of the host galaxies and therefore implications for the progenitors, we plot in Figure 11 the cumulative distribution of 80% light-radii (R_{80}) of our host galaxies. The median size for the LGRB sam-

ple is 3.0 kpc and the distribution spans from $R_{80} \sim 0.6$ kpc to $R_{80} \sim 10$ kpc. For comparison, we show the distributions for the samples of LGRBs and CCSNe studied by Svensson et al. (2010). A KS test reveals that our distribution, though shifted to slightly higher values of R_{80} , is consistent with their distribution of LGRB sizes. A KS test between our distribution of LGRB sizes and the CCSNe distribution yields a p-value of 3×10^{-3} indicating our larger sample of LGRBs is in full agreement with the conclusion of Svensson et al. (2010) that LGRB host galaxies are on average smaller than CCSNe host galaxies. We also show in Figure 11 the sample of SLSNe host galaxies studied by Lunnan et al. (2015). With the previous small sample of LGRBs, the SLSNe and LGRB host size distributions were statistically consistent. Now with our larger sample we find that the apparent larger sizes of LGRB hosts is statistically robust (KS p-value = 1.4×10^{-3}).

4.4. Durations and Energies

The durations (as measured by T_{90}) and isotropic equivalent energies ($E_{\gamma, \text{iso}}$) of LGRBs each vary by over several orders of magnitude. Whether or not environmental factors influence these variations in burst properties is an open question that we can address with our sample of offsets and fractional fluxes. In Figure 12 we plot fractional flux and host-normalized offset versus T_{90} and $E_{\gamma, \text{iso}}$. We find no significant correlations between fractional flux or offset with T_{90} or $E_{\gamma, \text{iso}}$, indicating that the locations of LGRBs within their host galaxies do not affect the properties of the bursts themselves. Our findings do not support the claim of a possible trend between host-normalized offset and isotropic equivalent energy reported by Ramirez-Ruiz et al. (2002) based on a much smaller sample of 16 pre-*Swift* events.

5. DISCUSSION

The distributions of offsets and fractional flux presented in Section 4 provide the most in-depth look at the locations and environments of LGRBs presented to date. Furthermore, our sample is sufficiently large and wide in redshift to allow investigation of trends with cosmic time. Here we discuss possible redshift evolution, the relationship between fractional flux and host-normalized offset, and the implications for the progenitors of LGRBs provided by this new view of their locations.

5.1. Redshift Trends

In Figure 13 we show the physical offsets for bursts with $\sigma_{R_{\text{phys}}} \lesssim 1.0$ kpc as a function of redshift (right panel) and the cumulative distribution of physical offsets binned into three equally populated bins of redshift (left panel): $z \leq 0.9$, $0.9 < z \leq 1.8$, and $z > 1.8$. We find that there is no statistically significant trend of the physical offsets as a function of redshift. KS tests between the three redshift bins yield p-values of 0.97, 0.77, and 0.45 for the low- z and mid- z bins, low- z and high- z bins, and mid- z and high- z bins, respectively.

In Figure 14 we show the host-normalized offsets of bursts with $\sigma_{R_{\text{norm}}} \lesssim 0.5$ as a function of redshift (right panel) and the cumulative distribution of host-normalized offsets binned by redshift (left panel). Overall we find no significant trend of the host-normalized offsets with redshift. KS tests between the three redshift bins yield p-values of 0.61, 0.51, and 0.23 for the low- z and mid- z bins, low- z and high- z bins, and mid- z and high- z bins, respectively. Despite the statistical insignificance, there appears to be a slight shift of the $z > 1.8$ bin toward larger offsets. If this reflects an actual trend, coupled




Figure 10. *Left:* Cumulative distributions of fractional flux for our sample (red), the LGRB sample of Fruchter et al. (2006) (F06, green), CCSNe from Svensson et al. (2010) (blue), SLSNe from Lunnan et al. (2015) (orange), and Type Ic (cyan) and II (violet) SNe from Kelly et al. (2008). We only include bursts in our sample with a ratio of error circle to galaxy area of $\lesssim 0.1$ (Figure 4). The diagonal dashed line shows the expected fractional flux distribution for a population uniformly tracing the underlying light of their host galaxies. *Right:* Cumulative distributions of fractional flux for two subsets of the sample divided at a rest-frame wavelength of 5000 Å. The agreement between the two means that we can use the rest-frame optical observations as a proxy for rest-frame UV.




Figure 11. Cumulative distributions of R_{80} for our sample of LGRBs (red), the LGRB (green) and CCSNe (blue) samples from Svensson et al. (2010) (S10), and the SLSNe sample from Lunnan et al. (2015) (orange). At high significance we find that LGRB host galaxy sizes are larger than SLSNe host sizes but smaller than CCSNe host sizes.

with the lack of a trend in the distribution of physical offsets, it would indicate an effect related to the sizes of the galaxies. In Figure 15 we show the measured half-light radii of the host galaxies as a function of redshift. There is clearly a trend of smaller galaxy sizes with increasing redshift. At

$z < 2$ the median galaxy half-light radius is 1.98 kpc and at $z > 2$ the median radius is 1.15 kpc. It is possible that surface brightness dimming, a strongly redshift dependent effect ($\propto (1+z)^{-4}$), may be limiting our ability to measure the sizes of the host galaxies at higher redshift. At higher redshifts the faint outer regions of galaxies become much harder to detect resulting in an underestimate of galaxy size and corresponding overestimate of the host normalized offset. However, *HST* observations of Lyman-break galaxies indicate that high redshift galaxies are intrinsically compact and show a size evolution of approximately $(1+z)^{-1}$ (Bouwens et al. 2004; Calvi et al. 2014). If low-mass galaxies such as LGRB hosts also lack diffuse outer components at high redshift, then it is possible the observed size trend in Figure 15 reflects actual galaxy growth.

In Figure 16 we show the fractional flux distribution binned into the same three redshift bins. KS tests between the three redshift bins yield p-values of 0.92, 0.13, and 0.42 for the low- z and mid- z bins, low- z and high- z bins, and mid- z and high- z bins, respectively. While the three binned distributions are statistically consistent, the lowest fractional flux values ($\lesssim 0.2$) only occur at $z > 1.5$. It is possible that surface brightness dimming is also playing a role here.

Overall we find no statistically significant trends with redshift in the offset or fractional flux distributions. The slight shift towards larger offsets and lower fractional flux in the highest redshift bin of $z > 1.8$ may indicate we are missing some of the diffuse outer parts of the hosts at high redshift.

5.2. The Fractional Flux - Offset Relationship

In Figure 17 we plot the fractional flux versus host-normalized offset for bursts satisfying error circle area to galaxy area ratio of $\lesssim 0.1$ and $\sigma_{R_{\text{norm}}} \lesssim 0.5$. We find a clear correlation between fractional flux and host-normalized off-




Figure 12. Scatter plots of fractional flux and host-normalized offset (R_{norm}) versus duration (T_{90}) and equivalent isotropic energy ($E_{\gamma,iso}$). We find no significant correlations between these quantities.




Figure 13. *Left:* Cumulative distribution of physical offsets for bursts with $\sigma_{R_{phys}} \lesssim 1.0$ kpc binned into three equal-numbered bins of redshift: $z \leq 0.9$ (blue), $0.9 < z \leq 1.8$ (red), and $z > 1.8$ (green). *Right:* Physical offsets with their associated uncertainties as a function of redshift. In both plots we do not include bursts with unknown redshift. We do not find a statistically significant trend in physical offset as a function of redshift.




Figure 14. Same as Figure 13 but for the host-normalized offsets of bursts with $\sigma_{R_{\text{norm}}} \lesssim 0.5$. We do not find a statistically significant trend in host-normalized offset as a function of redshift. The slight inclination towards larger offsets at higher redshift may be the result of surface brightness dimming.




Figure 15. Half-light radii as a function of redshift showing a clear trend that may explain the slight shift to larger host-normalized offsets at $z \gtrsim 2$. Whether this trend reflects real galaxy growth or the effects of surface brightness dimming is less clear.

set, where bursts at smaller offsets have high fractional flux values and bursts at larger offsets have on average lower fractional flux values. The scatter in the fractional flux values increases with increasing R_{norm} . All bursts in our sample with $R_{\text{norm}} \lesssim 0.5$ have fractional flux values greater than 0.6, yielding a notable lack of bursts with small offsets and low frac-

tional flux values. Beyond $R_{\text{norm}} \approx 0.5$ the fractional flux values are on average lower but also show considerable scatter. Bursts with $R_{\text{norm}} \gtrsim 1$ have fractional flux values spanning the full range from 0–1.

While it is not surprising that a trend should exist between fractional flux and offset given that regions close to the centers of galaxies are brighter, the coupling of fractional flux and offset information explains the results seen in the fractional flux distribution. To illustrate this, we also show in Figure 17 the cumulative distributions of fractional flux for the subsamples with $R_{\text{norm}} \leq 0.5$ and $R_{\text{norm}} > 0.5$. Bursts with $R_{\text{norm}} \leq 0.5$ exclusively occur on regions of high fractional flux and bursts with $R_{\text{norm}} > 0.5$ uniformly trace the light of their hosts, showing no preference for unusually bright regions. This means that the shift to high fractional flux we find in the total distribution is entirely due to LGRBs at small offsets and not to bursts at large offset that happen to reside in bright star forming regions. Therefore, it is not simply the presence of a bright star forming region that affects LGRB production, but rather another factor playing a role near the centers of galaxies must be at play.

Previous studies indicated that the agreement between LGRB offsets and the exponential disk was inconsistent with the observation that LGRBs were highly correlated with the brightest regions of their hosts (Bloom et al. 2002; Fruchter et al. 2006). The offset distribution we measure coupled with the cumulative distribution of fractional flux showing a preference for bright regions does not imply a discrepancy. Indeed, the fact that we find LGRBs concentrated in the inner regions of their galaxies with corresponding high fractional fluxes can be interpreted as a relief of the tension between the conclusions of Bloom et al. (2002) and Fruchter et al. (2006). Ultimately, the slight difference between our distribution of fractional flux and the distribution of Fruchter et al. (2006) may be explained by a difference in the offset distributions of




Figure 16. Same as Figure 13 but for fractional flux. We do not find a statistically significant trend in fractional flux as a function of redshift.




Figure 17. *Left:* Fractional flux versus host-normalized offset for LGRBs (red) with $\sigma_{R_{\text{norm}}} \lesssim 0.5$ and error circle to galaxy area ratio $\lesssim 0.1$ and Type Ic/Ic-BL SNe (cyan) from Kelly et al. (2008) and Kelly & Kirshner (2012). *Right:* Cumulative distributions of fractional flux for two subsets of the LGRB sample divided at $R_{\text{norm}} = 0.5$. There is a clear trend relating these two quantities where LGRBs at small host-normalized offset, near the bright cores of their hosts, are exclusively at high fractional flux. At $R_{\text{norm}} \gtrsim 0.5$, the bursts uniformly trace the underlying light of their host galaxies similar to CCSNe as also shown in the right panel. Therefore, the shift to high fractional flux seen in the full sample distribution (Figure 10) is due to bursts occurring in the inner parts of their host galaxies. A similar trend is seen in the Ic SNe sample.

the two samples, and may simply be the result of small number statistics where our sample more closely approximates the true underlying distribution.

In Figure 17 we also plot the normalized offsets and fractional fluxes of Type Ic and Ic-BL SNe for objects common to both the sample with fractional fluxes from Kelly et al. (2008)

and the sample with host-normalized offsets from Kelly & Kirshner (2012). We find a similar trend between the two quantities that indicates the shift to high fractional flux in the Ic SNe distribution can be attributed to SNe that occurred at small offset.

5.3. Progenitor Implications

We find that LGRBs are on average located at smaller host-normalized offset than expected for an exponential disk profile of star formation and are correspondingly more concentrated on brighter regions of their hosts than expected for a population simply tracing the underlying light distributions. The evidence presented here suggests that the progenitors of LGRBs prefer the central regions of their hosts, indicating that the properties of star formation towards the inner regions of LGRB host galaxies are favorable for LGRB production. As demonstrated in Figure 17, the bursts responsible for the strong preference towards higher fractional flux values are dominated by bursts at small offsets. The fractional flux distribution of bursts at offsets $R_{\text{norm}} \gtrsim 0.5$ uniformly tracks the host light. In effect the high fractional flux values are due to the fact that the bursts occur at small offsets where the host is brighter.

How the suspected metallicity bias of LGRB production fits into this new view of their preferred locations is an important question. Star forming disk galaxies typically have metallicity gradients with higher metallicities towards the cores (Vila-Costas & Edmunds 1992; Zaritsky et al. 1994). One might then expect, if metallicity gradients are also present in the hosts of LGRBs, to more frequently find LGRBs in the outskirts of their hosts contrary to what we find here. Recent resolved host studies of low redshift hosts indicate weak metallicity gradients (Levesque et al. 2011; Thöne et al. 2014) and that the global host metallicity is often a sufficient proxy for the metallicity of the burst site. Though the metallicity typically shows only slight variation across the host, the burst site often has the lowest metallicity (Christensen et al. 2008; Levesque et al. 2011; Thöne et al. 2014). In future work, we plan to test for the presence of metallicity gradients, in an average sense, in high redshift hosts by combining ISM metallicity measurements with our measured offsets. While it is clear that metallicity is a factor affecting the production of LGRBs, our results indicate that other factors relating to the mode of star formation may also play a dominant role. The conditions in the central regions of LGRB hosts, perhaps a consequence of their formation histories, may be such that the IMF is different, producing more massive stars and thus more potential LGRB progenitors. It has been shown that massive stars are often found in binary systems (Mason et al. 2009; Sana et al. 2012) and that binary interactions may be important for LGRB production (Fryer et al. 1999; Bromm & Loeb 2006). There may be increased numbers of massive star binaries in the central regions of LGRB hosts.

In both the offset and fractional flux distributions we find that LGRBs and Type Ic SNe prefer similar locations, indicating a common link between their progenitors. While LGRBs and Type Ic SNe broadly prefer similar locations, the exclusive association of LGRBs with Ic-BL SNe indicates that some factor must determine whether a star explodes as a normal Type Ic, a Type Ic-BL without a LGRB, or a Type Ic-BL with a LGRB. Previous observations suggest that the environments where Type Ic-BL SNe occur are more metal poor than those for normal Type Ic SNe, and that the environments of Type Ic-BL SNe associated with LGRBs are even more metal

poor (Modjaz et al. 2008, 2011; Kelly & Kirshner 2012). Recent work suggests that metallicity is likely not the only factor influencing the production of Ic-BL SNe and LGRBs. Kelly et al. (2014) found that Ic-BL SNe and LGRBs occur in host galaxies that have higher stellar mass and star formation rate densities than a control sample of SDSS star forming galaxies. They show that metallicity cannot explain this preference and instead suggest that another environmental factor such as increased numbers of young, bound clusters where tight binaries form or a top-heavy IMF may be at play in Ic-BL SNe and LGRB hosts. The results we find, namely that LGRBs prefer the central regions of their hosts, may be a sub-galactic manifestation of the same environmental factor responsible for the preference for high global stellar mass and star formation rate densities seen by Kelly et al. (2014). Whatever this factor is, it seems to influence primarily where LGRBs occur but not the properties of the explosions themselves as we do not find correlations between fractional flux or offset with burst duration or isotropic equivalent energy.

Although the SLSN sample size is much smaller than the LGRB sample size, the statistical consistency between their distributions of fractional flux may be hinting that SLSNe and LGRBs share a common mass progenitor. It could be that in the case of SLSNe, the core is not quite massive enough to form a black hole and instead collapses into a magnetar, while in the case of LGRBs, the core collapses into a black hole (Lunnan et al. 2014). However, the difference in the size distributions of SLSN and LGRB hosts indicates other factors also influence the production of their progenitors.

6. CONCLUSIONS

We have carried out a comprehensive study of the locations of LGRBs within their host galaxies using *HST* observations of *Swift* bursts compiled over the last decade. Using astrometry from ground- and space-based detections of optical, NIR, radio, and X-ray afterglows, we measure the projected physical and host-normalized offsets of 71 LGRBs from their host centers. In addition, we measure the fractional flux, the brightness of the burst site relative to the total host light distribution. Upon restricting the sample to avoid biases associated with measurement uncertainty we obtain sample sizes of 58, 52, and 50 LGRBs for the physical offsets, host-normalized offsets, and fractional fluxes, respectively. These measurements have enabled us to study the offset and fractional flux distributions in great detail, providing an in-depth view into the preferred locations of LGRBs. Our results are as follows:

- In agreement with previous work (Wainwright et al. 2007), we find that LGRB hosts typically show irregular morphologies and are on average smaller (median = 3.0 kpc) than the host galaxies of CCSNe, though not as small as the hosts of SLSNe.
- We find that the physical offsets of LGRBs have a mean of the Monte Carlo medians of 1.34 kpc with a 90% confidence interval of 1.17 – 1.51 kpc.
- The host-normalized offsets span a wide range of 0.04 to 7.0 but are concentrated at small values with a mean of the Monte Carlo medians of 0.67, with a 90% confidence interval of 0.58 – 0.76. This distribution of LGRB offsets is considerably more concentrated than expected if LGRBs traced an exponential disk (p-value = 2.6×10^{-4}). LGRBs are more concentrated than their

host galaxies' own light distributions, such that 50% of bursts reside within a region containing only 33% of the underlying host light.

- Compared to Type Ib/c SNe, LGRBs have consistent host-normalized offset distributions with 80% of synthetic Monte Carlo distributions yielding p-values > 0.05 . On the other hand, LGRB offsets are inconsistent with those of Type II SNe.
- The fractional flux distribution of our sample of LGRBs has a median at 0.75 and $\sim 80\%$ of bursts have fractional flux values $\gtrsim 0.5$, indicating a strong preference towards high fractional flux though not as strong as previous studies of smaller samples indicated.
- The host-normalized offsets and fractional fluxes are correlated, such that bursts at small offset ($\lesssim 0.5$) occur exclusively on regions of high fractional flux ($\gtrsim 0.6$) and bursts at larger offset uniformly track the underlying host light distributions, indicating that the preference for high fractional flux present in the full sample is entirely due to LGRBs at small offset.
- The offset and fractional flux distributions show no statistically significant trends with redshift, though a slight shift towards larger offsets and lower fractional flux values at $z \gtrsim 1.8$ may indicate the effects of surface brightness dimming.

From our study it appears that the star formation occurring within $\sim R_h$ is the most favorable for LGRB production. The distributions of offset and fractional flux for our sample of LGRBs agree well with those for Type Ic SNe, a scenario consistent with the well-known connection between LGRBs and Type Ic-BL SNe. Both types of transients occur in special types of galaxies and in special environments within their hosts, where metallicity is likely not the only important factor. In the central regions of galaxies it is possible that changes in the IMF or increased massive star binary fractions may be responsible for the sub-galactic preferences of LGRBs and Ic-BL SNe. This new view of LGRB locations we have presented is unlikely to change in the near future because the sample size is unlikely to increase significantly. Progress toward understanding the precise factors at play in the central regions of LGRB host galaxies will likely come through studies of LGRB progenitor analogs in the local universe whose environments can be studied in great detail.

The Berger GRB group at Harvard is supported in part by the NSF under grant AST-1411763 and by NASA under grant NNX15AE50G. This paper is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE1144152. W.F. acknowledges support provided by NASA through Einstein Postdoctoral Fellowship grant number PF4-150121 issued by the *Chandra* X-ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory for NASA under contract NAS8-03060. We thank Ragnhild Lunnan, Maria Drout, Josh Grindlay, and Avi Loeb for helpful discussions. The *HST* data presented here and the *Swift*/UVOT data used in this work were obtained from the Mikulski Archive for Space Telescopes (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA

contract NAS5-26555. Support for MAST for non-*HST* data is provided by the NASA Office of Space Science via grant NNX09AF08G and by other grants and contracts. Based in part on observations obtained at the Gemini Observatory, acquired through the Gemini Science Archive and processed using the Gemini IRAF package, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), Ministério da Ciência, Tecnologia e Inovação (Brazil) and Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina). Based in part on data obtained from the ESO Science Archive Facility. This paper makes use of data gathered with the 6.5 meter Magellan Telescopes located at Las Campanas Observatory, Chile. This research has made use of data obtained from the *Chandra* Data Archive and software provided by the *Chandra* X-ray Center (CXC) in the application package CIAO. Based in part on data collected at the Subaru Telescope and obtained from SMOKA, which is operated by the Astronomy Data Center, National Astronomical Observatory of Japan. We thank Brad Cenko for providing Palomar 60inch afterglow images for GRBs 060206 and 090618.

REFERENCES

- Alard, C. 2000, *A&AS*, 144, 363
 Beckwith, S. V. W., Stiavelli, M., Koekemoer, A. M., et al. 2006, *AJ*, 132, 1729
 Berger, E. 2010, *ApJ*, 722, 1946
 Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
 Bloom, J. S., Kulkarni, S. R., & Djorgovski, S. G. 2002, *AJ*, 123, 1111
 Bouwens, R. J., Illingworth, G. D., Blakeslee, J. P., Broadhurst, T. J., & Franx, M. 2004, *ApJ*, 611, L1
 Bouwens, R. J., Illingworth, G. D., Oesch, P. A., et al. 2015, *ApJ*, 803, 34
 Bromm, V., & Loeb, A. 2006, *ApJ*, 642, 382
 Calvi, V., Stiavelli, M., Bradley, L., Pizzella, A., & Kim, S. 2014, *ApJ*, 796, 102
 Christensen, L., Hjorth, J., & Gorosabel, J. 2004, *A&A*, 425, 913
 Christensen, L., Vreeswijk, P. M., Sollerman, J., et al. 2008, *A&A*, 490, 45
 Faber, S. M., Willmer, C. N. A., Wolf, C., et al. 2007, *ApJ*, 665, 265
 Frail, D. A., Kulkarni, S. R., Sari, R., et al. 2001, *ApJ*, 562, L55
 Fruchter, A. S., Levan, A. J., Strolger, L., et al. 2006, *Nature*, 441, 463
 Fryer, C. L., Woosley, S. E., & Hartmann, D. H. 1999, *ApJ*, 526, 152
 Gonzaga, S., & et al. 2012, *The DrizzlePac Handbook*, HST Data Handbook, Graham, J. F., & Fruchter, A. S. 2013, *ApJ*, 774, 119
 Hogg, D. W., Pahre, M. A., McCarthy, J. K., et al. 1997, *MNRAS*, 288, 404
 Kelly, P. L., Kirshner, R. P., & Pahre, M. 2008, *ApJ*, 687, 1201
 Kelly, P. L., & Kirshner, R. P. 2012, *ApJ*, 759, 107
 Kelly, P. L., Filippenko, A. V., Modjaz, M., & Kocevski, D. 2014, *ApJ*, 789, 23
 Krühler, T., Greiner, J., Schady, P., et al. 2011, *A&A*, 534, A108
 Le Floc'h, E., Duc, P.-A., Mirabel, I. F., et al. 2003, *A&A*, 400, 499
 Levesque, E. M., Kewley, L. J., Graham, J. F., & Fruchter, A. S. 2010a, *ApJ*, 712, L26
 Levesque, E. M., Kewley, L. J., Berger, E., & Zahid, H. J. 2010b, *AJ*, 140, 1557
 Levesque, E. M., Berger, E., Soderberg, A. M., & Chornock, R. 2011, *ApJ*, 739, 23
 Levesque, E. M. 2012, *Death of Massive Stars: Supernovae and Gamma-Ray Bursts*, 279, 167
 Lunnan, R., Chornock, R., Berger, E., et al. 2014, *ApJ*, 787, 138
 Lunnan, R., Chornock, R., Berger, E., et al. 2015, *ApJ*, 804, 90
 MacFadyen, A. I., & Woosley, S. E. 1999, *ApJ*, 524, 262
 Marchesini, D., van Dokkum, P., Quadri, R., et al. 2007, *ApJ*, 656, 42
 Mason, B. D., Hartkopf, W. I., Gies, D. R., Henry, T. J., & Helsel, J. W. 2009, *AJ*, 137, 3358
 Metcalfe, N., Shanks, T., Weilbacher, P. M., et al. 2006, *MNRAS*, 370, 1257
 Modjaz, M., Kewley, L., Kirshner, R. P., et al. 2008, *AJ*, 135, 1136
 Modjaz, M., Kewley, L., Bloom, J. S., et al. 2011, *ApJ*, 731, L4

- Ofek, E. O., Cenko, S. B., Gal-Yam, A., et al. 2007, *ApJ*, 662, 1129
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713
- Perley, D. A., Levan, A. J., Tanvir, N. R., et al. 2013, *ApJ*, 778, 128
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2014, *A&A*, 571, A16
- Podsiadlowski, P., Mazzali, P. A., Nomoto, K., Lazzati, D., & Cappellaro, E. 2004, *ApJ*, 607, L17
- Pollack, L. K., Chen, H.-W., Prochaska, J. X., & Bloom, J. S. 2009, *ApJ*, 701, 1605
- Prieto, J. L., Stanek, K. Z., & Beacom, J. F. 2008, *ApJ*, 673, 999
- Ramirez-Ruiz, E., Lazzati, D., & Blain, A. W. 2002, *ApJ*, 565, L9
- Sana, H., de Mink, S. E., de Koter, A., et al. 2012, *Science*, 337, 444
- Savaglio, S., Glazebrook, K., & Le Borgne, D. 2009, *ApJ*, 691, 182
- Schlafly, E. F., & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- Stanek, K. Z., Gnedin, O. Y., Beacom, J. F., et al. 2006, *AcA*, 56, 333
- Svensson, K. M., Levan, A. J., Tanvir, N. R., Fruchter, A. S., & Strolger, L.-G. 2010, *MNRAS*, 405, 57
- Thöne, C. C., Christensen, L., Prochaska, J. X., et al. 2014, *MNRAS*, 441, 2034
- van den Heuvel, E. P. J., & Yoon, S.-C. 2007, *Ap&SS*, 311, 177
- Vila-Costas, M. B., & Edmunds, M. G. 1992, *MNRAS*, 259, 121
- Vreeswijk, P., Malesani, D., Fynbo, J., et al. 2008, *GRB Coordinates Network*, 8301, 1
- Wainwright, C., Berger, E., & Penprase, B. E. 2007, *ApJ*, 657, 367
- Woosley, S. E. 1993, *ApJ*, 405, 273
- Woosley, S. E., & Bloom, J. S. 2006, *ARA&A*, 44, 507
- Zaritsky, D., Kennicutt, R. C., Jr., & Huchra, J. P. 1994, *ApJ*, 420, 87

Table 1
HST and Afterglow Observations

GRB	z	<i>HST</i> Image				Afterglow Image		
		Instrument	Filter	Exp. Time (s)	Obs. Date (UT)	Program ID	Telescope	Filter
040701	...	ACS/WFC1	F775W	1940	2004 Aug 09	10135	<i>Chandra</i>	...
040812	...	ACS/WFC1	F625W	1820				
			F775W	2120	2004 Oct 04	10135	<i>Chandra</i>	...
			F625W	2000				
040916	...	ACS/WFC1	F775W	2056	2004 Nov 30	10135
			F625W	1928				
040924	0.859	ACS/WFC1	F775W	3932	2005 Feb 18	10135	<i>HST</i> /ACS/WFC1	F775W
			F850LP	3932	2005 Feb 19			
041006	0.716	ACS/WFC1	F775W	4224	2005 Feb 10	10135	<i>HST</i> /ACS/WFC1	F775W
			F850LP	4224	2005 Feb 11			
050315	1.949	WFC3/IR	F160W	1209	2011 Jul 20	12307	Magellan/LDSS3	r
050401	2.9	WFC3/IR	F160W	1612	2010 Oct 1	12307	VLT/FORS2	R
050406	2.44	WFC3/IR	F160W	1612	2011 Feb 23	12307	Magellan/LDSS3	r
050408	1.236	WFC3/IR	F110W	1012	2013 Mar 24	12949	Gemini-N/GMOS	i
			F160W	1209				
050416A	0.654	ACS/WFC1	F775W	4224	2005 Nov 21	10135	<i>HST</i> /ACS/WFC1	F775W
			F850LP					
050525	0.606	ACS/WFC1	F625W	4268	2006 Mar 10	10135	<i>HST</i> /ACS/WFC1	F625W
			F775W					
050730	3.967	ACS/WFC1	F775W	7844	2010 Jun 10	11734	LCO/Swope	WashT1
050803	...	WFC3/IR	F160W	906	2011 Sep 3	12307	<i>Swift</i> /XRT	...
050820	2.612	ACS/WFC1	F850LP	14280	2006 Jun 8+11	10551	<i>HST</i> /ACS/WFC1	F850LP
			F625W	2238	2006 Jun 5			
			F775W	4404				
050824	0.83	WFC3/IR	F160W	906	2011 Jan 18	12307	VLT/FORS2	R
050904	6.29	ACS/WFC1	F850LP	4216	2005 Sep 26	10616	<i>HST</i> /NICMOS	F160W
		NICMOS	F160W	15359	2006 Jul 23			
050908	3.344	ACS/WFC1	F775W	7892	2010 Oct 31	11734	Gemini-S/GMOS	r
051016B	0.936	WFC3/IR	F160W	906	2011 Jan 16	12307	<i>Swift</i> /UVOT	U+B+V
051022	0.8	ACS/WFC1	F606W	1560	2009 Aug 21	11343	<i>Chandra</i>	...
		WFC3/IR	F160W	2397	2009 Oct 12			
060115	3.53	ACS/WFC1	F814W	7910	2010 Aug 27	11734	VLT/FORS1	I
060116	...	NICMOS	F160W	5120	2006 Dec 12	10633	<i>HST</i> /NICMOS	F160W
			F110W	5120	2006 Dec 11			
		ACS/WFC1	F775W	4400	2006 Oct 10			
			F850LP	1650	2006 Oct 11			
060124	2.296	WFC3/IR	F160W	1612	2010 Sep 28	12307	<i>Swift</i> /UVOT	V
060206	4.048	ACS/WFC1	F814W	9886	2006 Nov 25	10817	Palomar 60inch	I
060218	0.033	WFC3/IR	F160W	906	2010 Oct 12	12307	<i>HST</i> /ACS/WFC1	F625W
		ACS/WFC1	F625W	2040	2006 Oct 26	10551		
			F814W	2040				
			F435W	1530	2006 Nov 04			
			F555W	1020				
060223	4.41	WFC3/IR	F110W	8395	2010 Sep 13	11734	<i>Swift</i> /UVOT	V
060418 ^a	1.49	ACS/WFC1	F625W	4220	2006 May 09	10551	Self- <i>HST</i>	...
			F775W	4220				
			F555W	4386	2006 Jul 13			
060502A	1.51	WFC3/IR	F160W	1209	2010 Oct 11	12307	Gemini-N/GMOS	r
060505 ^b	0.089	ACS/WFC1	F475W	27774	2006 May+Jun	10551	Gemini-S/GMOS	g
			F814W	6840	2006 Jun 6+7			
		WFC3/IR	F160W	906	2011 Aug 3	12307		
060512 ^c	0.443	WFC3/IR	F160W	453	2011 Feb 23	12307	VLT/FORS1	...
060522	5.11	WFC3/IR	F110W	8395	2010 Oct 17	11734	<i>Swift</i> /UVOT	White
060526	3.21	ACS/WFC1	F775W	7844	2009 Aug 9	11734	VLT/FORS1	R
060602A	0.787	WFC3/IR	F160W	906	2010 Dec 5	12307	<i>Swift</i> /XRT	...
060605	3.78	ACS/WFC1	F775W	7862	2010 Oct 6	11734	<i>Swift</i> /UVOT	W1+V
060607	3.082	ACS/WFC1	F775W	7910	2010 Sep 17	11734	Gemini-N/GMOS	r
060614	0.125	ACS/WFC1	F606W	3600	2006 Sep 8	10917	<i>HST</i> /WFPC2	F606W
			F435W	2152				
			F814W	4840	2006 Oct 31			
		WFC3/IR	F160W	906	2010 Oct 8	12307		
060719	1.532	WFC3/IR	F160W	1209	2013 Feb 24	12949	VLT/FORS2	R
			F125W	1059				
060729	0.54	WFC3/IR	F160W	906	2010 Sep 15	12307	Gemini-S/GMOS	i
060912A	0.937	WFC3/IR	F160W	906	2011 Sep 23	12307	<i>Swift</i> /UVOT	White
060923A	...	WFC3/IR	F160W	1198	2013 Feb 10	12949	Gemini-N/NIRI	K
			F125W	1348				
060927	5.6	WFC3/IR	F110W	13992	2010 Sep 25	11734	VLT/FORS2	I
		NICMOS	F160W	10240	2007 Jun 29	10926		
061007	1.261	WFC3/IR	F160W	1209	2011 Jul 8	12307	VLT/FORS1	R
061110A	0.758	WFC3/IR	F160W	906	2010 Sep 30	12307	VLT/FORS1	R
061110B	3.44	ACS/WFC1	F775W	7862	2010 Sep 23	11734	VLT/FORS1	R
061222A	2.088	NICMOS	F160W	7680	2007 Jun 16	10908	Gemini-N/NIRI	K _s

Table 1 — *Continued*

GRB	z	HST Image				Afterglow Image		
		Instrument	Filter	Exp. Time (s)	Obs. Date (UT)	Program ID	Telescope	Filter
070125	1.547	WFC3/IR	F110W	2812	2010 Apr 24	11717	Gemini-N/GMOS	r
070208	1.165	WFC3/UVIS	F336W	2520	2013 Jan 02	12949	Gemini-N/GMOS	r
		WFC3/IR	F160W	1209				
070306	1.496	WFC3/IR	F110W	1012	2012 Nov 15	12949	VLT/ISAAC	K _s
			F160W	1198				
070318	0.836	WFC3/IR	F125W	1348	2010 Dec 31	12307	VLT/FORS1	R
			F160W	906				
070508	...	WFC3/IR	F160W	906	2011 Feb 22	12307	Magellan/MAGIC	I
070521	1.35	WFC3/IR	F160W	906	2011 Aug 02	12307	Swift/XRT	...
070721B	3.626	ACS/WFC1	F775W	7844	2010 Nov 13	11734	VLT/FORS2	R
070802	2.45	WFC3/IR	F160W	1209	2013 Feb 22	12949	VLT/FORS2	I
			F105W	1059				
071010A	0.98	WFC3/IR	F160W	906	2010 Oct 29	12307
071010B	0.947	WFC3/IR	F160W	906	2010 Nov 25	12307	Gemini-N/GMOS	r
071031	2.692	WFC3/IR	F160W	1612	2010 Nov 20	12307	VLT/FORS2	V
071112C	0.823	WFC3/IR	F160W	906	2010 Oct 08	12307	Gemini-N/GMOS	r
071122	1.14	WFC3/IR	F160W	1209	2010 Dec 21	12307	Gemini-N/GMOS	i
080207	2.086	WFC3/IR	F110W	2397	2009 Dec 09	11343	Chandra	...
		WFPC2	F606W	1600	2008 Mar 18			
			F814W	3320	2009 Mar 20			
			F702W	3600	2009 Mar 21			
		NICMOS	F160W	2560	2008 Apr 05			
		WFPC2	F606W	3200	2008 Jul 03			
080319B ^d	0.937		F814W	3200	2008 Jul 05	11513	HST/WFPC2	...
080319C	1.95	WFC3/IR	F160W	1209	2010 Sep 19	12307	Gemini-N/GMOS	r
080325	1.78	WFC3/IR	F160W	1209	2012 Dec 31	12949	Subaru/MOIRCS	K _s
			F125W	1359				
080430	0.767	WFC3/IR	F160W	906	2011 Jun 21	12307	Swift/UVOT	U+White
080520	1.545	WFC3/IR	F160W	1209	2011 Feb 08	12307	Swift/UVOT	White
080603A	1.688	WFC3/IR	F125W	1397	2013 Apr 10	12949	Gemini-N/GMOS	r
			F160W	1348				
080603B	2.69	WFC3/IR	F160W	1612	2011 Aug 06	12307	Swift/UVOT	U+B+V+W1
080605 ^b	1.640	WFC3/IR	F160W	2418	2012+2013	12307+12949	VLT/FORS2	R
			F125W	1209	2013 Mar 15			
080607	3.036	WFC3/IR	F160W	10791	2010 Jul 25	12005	Swift/UVOT	White+V+W2
080707	1.23	WFC3/IR	F160W	1209	2010 Oct 31	12307	VLT/FORS1	R
080710	0.845	WFC3/IR	F160W	906	2011 Feb 12	12307	Gemini-S/GMOS	r
080805	1.505	WFC3/IR	F160W	1209	2011 Oct 01	12307	VLT/FORS2	I
080913	6.7	WFC3/IR	F160W	7818	2009 Nov 30	11189	VLT/FORS2	z
080916A	0.689	WFC3/IR	F160W	906	2011 Mar 19	12307	VLT/FORS1	R
080928	1.692	WFC3/IR	F160W	1209	2010 Sep 18	12307	Gemini-S/GMOS	r
081007	0.530	WFC3/IR	F160W	906	2010 Nov 30	12307	Gemini-S/GMOS	r
081008	1.969	WFC3/IR	F160W	1209	2011 Sep 04	12307	Gemini-S/GMOS	r
081109	0.979	WFC3/IR	F160W	1359	2013 Mar 09	12949
			F110W	1312				
081121	2.512	WFC3/IR	F160W	1612	2011 Jan 04	12307	Swift/UVOT	U+B+V+White
081221	2.26	WFC3/IR	F160W	1209	2013 Jun 17	12949	Gemini-N/NIRI	K
			F105W	1209				
090113	1.749	WFC3/IR	F160W	2612	2009 Oct 17	11840
			ACS/WFC1	F606W	2208			
090404	...	WFC3/IR	F160W	2612	2010 Jan 09	11840	PdBI	108 GHz
			F105W	1359	2013 Jul 06			
			F125W	1209				
			F606W	2208				
090407	1.449	ACS/WFC1	F160W	1209	2010 Sep 02	11840	Chandra	...
		WFC3/IR	F160W	1209	2010 Sep 15	11840		
		WFC3/UVIS	F606W	740				
			F160W	2612				
090417B	0.345	WFC3/IR	F160W	2612	2009 Oct 17	11840	Swift/XRT	...
		ACS/WFC1	F606W	1656	2011 Jan 22			
			F160W	1209				
090418A	1.608	WFC3/IR	F160W	1209	2010 Oct 02	12307	Swift/UVOT	U
090423 ^b	8.2	WFC3/IR	F160W	42997	2010+2011	11189	Gemini-N/NIRI	H
			F125W	52117	2010			
090424	0.544	WFC3/IR	F160W	906	2011 May 03	12307	Gemini-S/GMOS	r
090429B ^b	9.4	WFC3/IR	F160W	4823	2010 Jan+Feb	11189	Gemini-N/NIRI	...
			F105W	4823	2010 Feb 24+28			
		ACS/WFC1	F606W	2100	2010 Jan 03			
			F160W	906				
090618	0.54	WFC3/IR	F160W	906	2011 Jan 09	12307	Palomar 60inch	I
090709A	...	WFC3/IR	F160W	1359	2012 Oct 15	12949	Subaru/MOIRCS	K _s
			F125W	1397				
091127	0.49	WFC3/IR	F160W	906	2010 Dec 16	12307	Gemini-S/GMOS	i
091208B	1.063	WFC3/IR	F160W	1209	2010 Oct 10	12307	Gemini-N/GMOS	r
100205A	...	WFC3/IR	F160W	1209	2010 Dec 06	11840	Gemini-N/NIRI	K
			F606W	1140				
100316D	0.059	WFC3/UVIS	F606W	1140	2011 Aug 03	12323	HST/WFC3/UVIS	F336W
		WFC3/UVIS	F336W	1680				
			F225W	2520	2011 Aug 02			

Table 1 — *Continued*

GRB	z	<i>HST</i> Image				Afterglow Image			
		Instrument	Filter	Exp. Time (s)	Obs. Date (UT)	Program ID	Telescope	Filter	
100413A	3.9	WFC3/IR	F438W	1230	2011 Aug 03	11840	EVLA	8.5 GHz	
			F814W	1260					
			F275W	2520					
			F555W	1680					
			F625W	720	2011 Aug 06				
			F763M	1720					
			F125W	1312	2011 Aug 02				
			F160W	1312					
			ACS/WFC1	FR716N	2400				2011 Aug 06
			WFC3/IR	F160W	1209				2010 Aug 31
WFC3/UVIS	F606W	752							
100526A	...	WFC3/IR	F125W	1348	2013 Jun 30	12949	Gemini-N/NIRI	K	
100615A	1.398	WFC3/UVIS	F606W	1128	2010 Dec 16	11840	<i>Chandra</i>	...	
		WFC3/IR	F160W	1209					
100621A	0.542	WFC3/IR	F160W	909	2013 Mar 09	12949	VLT/XSHOOTER	I	
			F105W	748					
100905A	...	WFC3/IR	F140W	5440	2013 Jul 10	12247	
110312A	...	WFC3/IR	F160W	1209	2011 Nov 18	12378	<i>Swift</i> /XRT	...	
		WFC3/UVIS	F606W	1110	2011 Nov 17				
110709B	...	WFC3/UVIS	F606W	2480	2011 Nov 08	12378	<i>Chandra</i>	...	
		WFC3/IR	F160W	2612	2011 Nov 12				
110731A	2.83	WFC3/UVIS	F625W	5200	2012 Aug 22	12370	<i>HST</i> /WFC3/UVIS	F625W	
120119A	1.728	WFC3/IR	F160W	1209	2012 Oct 08	12949	Gemini-S/GMOS	r	
			F125W	1209					

Note. — The table is organized such that the first image listed for each burst is the image for which the corresponding offset and fractional flux measurements were used to compile the distribution (Section 4). For bursts with multiple observations we group the observations by instrument. Within an instrument group, the observations are listed by date. Only the last epoch of observations are shown for bursts with multiple epochs, unless an earlier epoch was chosen as the best host image. Blank entries in the table indicate inheritance of the entry in the previous row.

^a The afterglow and what we believe to be the host galaxy (Pollack et al. 2009) are both detected in the same epoch and offset from each other such that centroid measurements are not biased. Therefore the offset (see Section 3.1) can be measured using this single early epoch image. To measure the fractional flux (see Section 3.4) we used a late epoch image where the afterglow has faded away.

^b We combined multiple epochs in the same filter to obtain deep images. For GRB 060505, the afterglow was not detected in early epoch *HST* observations so we combined these observations with a later epoch.

^c The *HST* data for GRB 060512 is of poor quality, preventing further analysis.

^d We combined the F606W and F814W filters to obtain a higher S/N detection of the host. The afterglow image is also a stack of these two filters at an earlier epoch.

Table 2
Offsets, Galaxy Sizes, Magnitudes, and P_{cc}

GRB	# Tie Objects	σ_{tie} ($''$)	σ_{AG} ($''$)	σ_{host} ($''$)	R ($''$)	R_{phys} (kpc)	R_h ($''$)	R_{norm}	AB Mag ^a	P_{cc}
040701	4	0.367	0.475
040812	5	0.115	0.046	0.019	0.270 ± 0.125	2.283 ± 1.057	0.297	0.908 ± 0.420	19.450 ± 0.010	<0.001
040916	0.150	>27.1	...
040924	41	0.005	0.002	0.014	0.286 ± 0.015	2.197 ± 0.116	0.188	1.521 ± 0.080	23.954 ± 0.024	0.003
041006 ^b	24	0.004	<0.001	0.007	0.352 ± 0.008	2.536 ± 0.061	0.298	1.180 ± 0.029	24.668 ± 0.057	0.012
050315	17	0.054	0.003	0.017	0.114 ± 0.056	0.953 ± 0.474	0.236	0.481 ± 0.239	23.808 ± 0.029	0.006
050401	24	0.044	0.025	0.009	0.082 ± 0.051	0.638 ± 0.397	0.170	0.483 ± 0.300	25.217 ± 0.077	0.006
050406	7	0.038	0.217	0.006	0.195 ± 0.220	1.579 ± 1.788	0.121	1.609 ± 1.822	26.602 ± 0.140	0.041
050408	14	0.027	0.002	0.010	0.152 ± 0.029	1.263 ± 0.239	0.278	0.545 ± 0.103	23.700 ± 0.022	0.008
050416A	24	0.007	<0.001	0.011	0.037 ± 0.013	0.260 ± 0.090	0.160	0.234 ± 0.081	23.001 ± 0.009	0.001
050525	74	0.006	0.001	0.002	0.028 ± 0.007	0.185 ± 0.046	0.119	0.232 ± 0.057	26.067 ± 0.064	0.005
050730	9	0.038	0.014	>27.9	...
050803	0.850
050820	28	0.009	0.003	0.004	0.442 ± 0.010	3.530 ± 0.081	0.116	3.808 ± 0.087	25.196 ± 0.050	0.010
050824	6	0.036	0.012	0.038	0.473 ± 0.054	3.597 ± 0.410	0.402	1.177 ± 0.134	23.716 ± 0.042	0.021
050904	9	0.027	0.020	>27.2	...
050908	15	0.054	0.001	0.003	0.037 ± 0.055	0.275 ± 0.405	0.068	0.544 ± 0.802	27.669 ± 0.136	0.005
051016B	4	0.062	0.184	0.013	0.257 ± 0.194	2.023 ± 1.532	0.228	1.127 ± 0.853	22.314 ± 0.011	0.004
051022	5	0.121	0.015	0.021	0.171 ± 0.124	1.283 ± 0.932	0.334	0.512 ± 0.372	22.020 ± 0.006	0.002
060115	18	0.045	0.043	0.017	0.280 ± 0.065	2.042 ± 0.471	0.125	2.239 ± 0.517	27.342 ± 0.137	0.030
060116	10	0.024	0.029	>27.1	...
060124	7	0.058	0.065	0.025	0.132 ± 0.091	1.081 ± 0.745	0.172	0.766 ± 0.528	26.133 ± 0.126	0.011
060206	6	0.127	0.021	0.005	0.292 ± 0.128	2.021 ± 0.888	0.091	3.211 ± 1.411	27.647 ± 0.094	0.035
060218	13	0.012	<0.001	0.044	0.174 ± 0.045	0.115 ± 0.030	0.606	0.287 ± 0.075	19.635 ± 0.003	0.002
060223	9	0.173	0.041	0.009	0.112 ± 0.178	0.749 ± 1.186	0.211	0.532 ± 0.844	26.534 ± 0.069	0.026
060418	0.003	0.004	0.404 ± 0.005	3.419 ± 0.040	0.118	3.425 ± 0.040	25.855 ± 0.067	0.019
060502A	8	0.021	0.009	0.006	0.051 ± 0.023	0.435 ± 0.198	0.162	0.317 ± 0.144	25.864 ± 0.110	0.007
060505	20	0.029	0.004	0.023	4.307 ± 0.037	7.160 ± 0.062	1.195	3.605 ± 0.031	19.101 ± 0.001	0.009
060522	4	0.089	0.146	>28.9	...
060526	23	0.040	0.006	>27.4	...
060602A	2.550
060605	12	0.115	0.020	0.002	0.183 ± 0.117	1.304 ± 0.829	0.109	1.683 ± 1.069	27.555 ± 0.148	0.026
060607	33	0.037	0.003	>28.9	...
060614	21	0.009	0.002	0.011	0.357 ± 0.014	0.801 ± 0.031	0.329	1.086 ± 0.042	22.821 ± 0.007	0.003
060719	12	0.051	0.062	0.011	0.206 ± 0.081	1.743 ± 0.684	0.246	0.837 ± 0.328	23.400 ± 0.023	0.006
060729	17	0.039	0.001	0.015	0.338 ± 0.042	2.149 ± 0.265	0.308	1.098 ± 0.135	23.509 ± 0.028	0.011
060912A	11	0.148	0.005	0.036	0.654 ± 0.153	5.155 ± 1.203	0.713	0.918 ± 0.214	21.547 ± 0.013	0.017
060923A	38	0.039	0.013	0.052	0.291 ± 0.066	2.460 ± 0.560	0.696	0.418 ± 0.095	23.278 ± 0.038	0.038
060927	18	0.026	0.111	>28.6	...
061007	5	0.026	0.100	<0.001	0.022 ± 0.103	0.184 ± 0.862	0.373	0.059 ± 0.277	23.621 ± 0.033	0.013
061110A	11	0.035	0.014	0.015	0.135 ± 0.040	0.992 ± 0.297	0.176	0.765 ± 0.229	25.114 ± 0.083	0.007
061110B	49	0.025	0.011	0.011	0.061 ± 0.030	0.449 ± 0.220	0.104	0.586 ± 0.288	27.230 ± 0.105	0.008
061222A	15	0.026	0.034	0.011	0.044 ± 0.044	0.368 ± 0.368	0.193	0.229 ± 0.229	24.399 ± 0.050	0.005
070125	5	0.033	0.007	>28.0	...
070208	6	0.030	0.006	0.025	0.099 ± 0.040	0.820 ± 0.327	0.368	0.270 ± 0.108	21.902 ± 0.009	0.005
070306	14	0.043	0.005	0.011	0.090 ± 0.045	0.760 ± 0.380	0.209	0.430 ± 0.215	21.860 ± 0.007	0.002
070318	8	0.057	0.007	0.019	0.109 ± 0.060	0.827 ± 0.458	0.205	0.529 ± 0.293	24.100 ± 0.034	0.005
070508	16	0.025	0.004	0.020	0.423 ± 0.033	3.577 ± 0.277	0.592	0.714 ± 0.055	22.871 ± 0.031	0.027
070521	0.850
070721B	17	0.042	0.032	0.006	0.045 ± 0.054	0.327 ± 0.387	0.057	0.793 ± 0.939	28.442 ± 0.165	0.011
070802	10	0.039	0.007	0.027	0.132 ± 0.048	1.069 ± 0.390	0.348	0.379 ± 0.138	23.994 ± 0.043	0.014
071010A	11	0.127	0.300
071010B	7	0.022	0.001	0.010	0.109 ± 0.025	0.860 ± 0.195	0.257	0.424 ± 0.096	22.635 ± 0.016	0.004
071031	7	0.068	<0.001	>27.4	...
071112C	10	0.020	0.007	0.017	0.202 ± 0.027	1.533 ± 0.206	0.398	0.508 ± 0.068	23.674 ± 0.046	0.016
071122	10	0.032	0.033	0.025	0.075 ± 0.053	0.616 ± 0.433	0.393	0.191 ± 0.134	22.678 ± 0.017	0.009
080207	2	0.381	0.206	0.066	0.769 ± 0.438	6.402 ± 3.650	0.683	1.126 ± 0.642	23.225 ± 0.025	0.047
080319B	18	0.011	0.001	0.117	0.215 ± 0.118	1.697 ± 0.929	0.147	1.465 ± 0.802	26.679 ± 0.200	0.015
080319C	10	0.018	0.008	0.046	0.855 ± 0.050	7.177 ± 0.417	0.638	1.341 ± 0.078	22.018 ± 0.011	0.023
080325	60	0.022	0.041	0.059	0.697 ± 0.075	5.890 ± 0.635	0.443	1.574 ± 0.170	22.537 ± 0.016	0.017
080430	4	0.035	0.015	0.030	0.108 ± 0.049	0.800 ± 0.361	0.282	0.384 ± 0.173	24.595 ± 0.071	0.011
080520	17	0.106	0.146	0.024	0.468 ± 0.182	3.966 ± 1.543	0.223	2.100 ± 0.817	23.965 ± 0.032	0.011
080603A	22	0.027	<0.001	0.013	0.089 ± 0.030	0.753 ± 0.255	0.202	0.440 ± 0.149	22.884 ± 0.011	0.003
080603B	6	0.117	0.034	>26.7	...
080605	38	0.029	0.009	0.008	0.081 ± 0.031	0.682 ± 0.266	0.198	0.407 ± 0.159	22.467 ± 0.008	0.002
080607	5	0.230	0.056	0.024	0.681 ± 0.238	5.229 ± 1.827	0.419	1.626 ± 0.568	24.372 ± 0.027	0.036
080707	6	0.047	0.024	0.009	0.089 ± 0.054	0.739 ± 0.449	0.263	0.338 ± 0.205	22.950 ± 0.017	0.005
080710	6	0.035	<0.001	>26.1	...
080805	15	0.038	0.006	0.026	0.466 ± 0.047	3.944 ± 0.396	0.275	1.695 ± 0.170	23.249 ± 0.020	0.010
080913	13	0.035	0.044	>28.0	...
080916A	12	0.056	0.022	0.010	0.011 ± 0.061	0.075 ± 0.432	0.240	0.044 ± 0.254	22.762 ± 0.016	0.004
080928	16	0.021	0.012	0.016	1.705 ± 0.029	14.437 ± 0.245	0.442	3.858 ± 0.066	21.990 ± 0.010	0.035
081007	11	0.023	<0.001	0.017	0.111 ± 0.029	0.700 ± 0.182	0.277	0.402 ± 0.105	24.489 ± 0.059	0.011
081008	28	0.021	0.002	0.018	1.658 ± 0.028	13.902 ± 0.236	0.236	7.027 ± 0.119	23.591 ± 0.028	0.066

Table 2 — *Continued*

GRB	# Tie Objects	σ_{tie} ($''$)	σ_{AG} ($''$)	σ_{host} ($''$)	R ($''$)	R_{phys} (kpc)	R_h ($''$)	R_{norm}	AB Mag ^a	P_{cc}
081109	0.850
081121	6	0.12	0.014	0.011	0.262 ± 0.122	2.112 ± 0.981	0.198	1.323 ± 0.615	24.923 ± 0.054	0.010
081221	11	0.047	0.016	0.055	0.388 ± 0.074	3.195 ± 0.609	0.409	0.949 ± 0.181	23.156 ± 0.025	0.013
090113	0.910
090404	4	0.115	0.361	0.044	0.556 ± 0.381	4.709 ± 3.229	0.690	0.806 ± 0.553	23.370 ± 0.018	0.045
090407	5	0.134	0.387	0.021	0.172 ± 0.410	1.451 ± 3.463	0.336	0.511 ± 1.220	22.953 ± 0.019	0.026
090417B	0.850
090418A	5	0.209	0.014	0.017	0.096 ± 0.211	0.812 ± 1.785	0.214	0.448 ± 0.984	23.687 ± 0.025	0.009
090423	6	0.031	0.005	>28.2	...
090424	9	0.021	0.005	0.030	0.407 ± 0.037	2.594 ± 0.235	0.407	1.000 ± 0.090	21.260 ± 0.007	0.005
090429B	10	0.053	0.012	>28.1	...
090618	12	0.038	0.001	0.025	0.693 ± 0.046	4.402 ± 0.292	0.519	1.335 ± 0.089	22.464 ± 0.020	0.020
090709A	25	0.028	0.021	0.007	0.283 ± 0.036	2.393 ± 0.305	0.256	1.105 ± 0.141	24.213 ± 0.038	0.010
091127	16	0.040	0.005	0.028	0.221 ± 0.049	1.335 ± 0.296	0.411	0.538 ± 0.119	22.700 ± 0.021	0.011
091208B	7	0.019	0.001	>26.7	...
100205A	12	0.034	0.028	>26.7	...
100316D ^c	15	0.007	<0.001	<0.001
100413A	8	0.098	0.600
100526A	10	0.047	0.008	0.008	0.194 ± 0.049	1.645 ± 0.411	0.188	1.034 ± 0.258	24.766 ± 0.033	0.007
100615A	3	0.318	0.028	0.004	0.246 ± 0.320	2.073 ± 2.694	0.083	2.963 ± 3.850	25.092 ± 0.033	0.034
100621A	6	0.062	0.025	0.015	0.045 ± 0.069	0.284 ± 0.439	0.262	0.17 ± 0.263	21.316 ± 0.006	0.002
100905A	0.910
110312A	0.850
110709B	1	0.318	0.057	0.005	0.067 ± 0.323	0.564 ± 2.737	0.073	0.912 ± 4.430	26.714 ± 0.080	0.111
110731A	211	0.006	0.001	0.008	0.182 ± 0.010	1.423 ± 0.082	0.121	1.502 ± 0.087	25.236 ± 0.043	0.005
120119A	49	0.013	<0.001	0.012	0.012 ± 0.017	0.099 ± 0.146	0.202	0.058 ± 0.086	23.480 ± 0.022	0.003

Note. — For bursts with multiple *HST* images of the host galaxy, the image used to make the measurements listed here corresponds to the first image listed for such bursts in Table 1. We show the 3σ upper limits for bursts with no host detection. Bursts with no listed offset or upper limit correspond to those bursts for which we were not able to obtain a suitable afterglow image.

^a Magnitudes are corrected for Galactic extinction (Schlafly & Finkbeiner 2011)

^b The SN associated with GRB 041006 is detected in the final epoch. The SN and host center are sufficiently offset from each other such that centroid measurements are not biased. However, measurement of fractional flux is not possible.

^c GRB 100316D occurred in a low-redshift galaxy with irregular morphology. The complexity of the host system prevents a meaningful definition of the host center. We therefore do not make an offset measurement for this burst.

Table 3
Fractional Flux and Ratio of Error Circle to Galaxy Area

GRB	Fractional Flux	Ratio	GRB	Fractional Flux	Ratio
040701	070802	0.692	1.30×10^{-2}
040812	0.423	1.74×10^{-1}	071010A
040916	071010B	0.825	7.34×10^{-3}
040924	0.783	8.21×10^{-4}	071031
041006	...	1.80×10^{-4}	071112C	0.774	2.83×10^{-3}
050315	0.693	5.25×10^{-2}	071122	0.951	1.37×10^{-2}
050401	0.799	8.86×10^{-2}	080207	0.207	4.02×10^{-1}
050406	0.110	3.31	080319B	0.598	5.65×10^{-3}
050408	0.724	9.48×10^{-3}	080319C	0.599	9.53×10^{-4}
050416A	0.896	1.91×10^{-3}	080325	0.134	1.10×10^{-2}
050525	0.914	2.61×10^{-3}	080430	0.645	1.82×10^{-2}
050730	080520	0.117	6.55×10^{-1}
050803	080603A	0.880	1.79×10^{-2}
050820	0.606	6.69×10^{-3}	080603B
050824	0.324	8.91×10^{-3}	080605	0.820	2.35×10^{-2}
050904	080607	0.177	3.19×10^{-1}
050908	0.316	6.31×10^{-1}	080707	0.836	4.03×10^{-2}
051016B	0.329	7.25×10^{-1}	080710
051022	0.556	1.33×10^{-1}	080805	0.517	1.96×10^{-2}
060115	0.081	2.48×10^{-1}	080913
060116	080916A	0.930	6.28×10^{-2}
060124	0.427	2.57×10^{-1}	080928	0.009	2.99×10^{-3}
060206	0.032	2.00	081007	0.792	6.89×10^{-3}
060218	0.872	3.92×10^{-4}	081008	0.000	7.99×10^{-3}
060223	0.292	7.10×10^{-1}	081109
060418	0.000	6.46×10^{-4}	081121	0.197	3.72×10^{-1}
060502A	0.689	1.99×10^{-2}	081221	0.761	1.47×10^{-2}
060505	0.987	6.00×10^{-4}	090113
060522	090404	0.183	3.02×10^{-1}
060526	090407	0.245	1.49
060602A	090417B
060605	0.103	1.15	090418A	0.367	9.58×10^{-1}
060607	090423
060614	0.459	7.85×10^{-4}	090424	0.740	2.81×10^{-3}
060719	0.581	1.07×10^{-1}	090429B
060729	0.246	1.60×10^{-2}	090618	0.260	5.36×10^{-3}
060912A	0.525	4.31×10^{-2}	090709A	0.461	1.87×10^{-2}
060923A	0.980	3.49×10^{-3}	091127	0.833	9.62×10^{-3}
060927	091208B
061007	0.778	7.67×10^{-2}	100205A
061110A	0.419	4.59×10^{-2}	100316D	0.879	...
061110B	0.393	6.90×10^{-2}	100413A
061222A	0.955	4.92×10^{-2}	100526A	0.531	6.43×10^{-2}
070125	100615A	0.051	14.8
070208	0.928	6.91×10^{-3}	100621A	0.914	6.51×10^{-2}
070306	0.798	4.29×10^{-2}	100905A
070318	0.781	7.85×10^{-2}	110312A
070508	0.517	1.83×10^{-3}	110709B	0.055	19.6
070521	110731A	0.029	2.53×10^{-3}
070721B	0.211	8.58×10^{-1}	120119A	1.000	4.14×10^{-3}

Table 4
Summary of KS Test p-values

LGRBs - This Work	Exp. Disk	Type Ib/c SNe ^a	Type Ic SNe	Type II SNe	CCSNe	SLSNe	LGRBs - F06	LGRBs - B02
Physical Offsets	...	1.33×10^{-6}	...	1.53×10^{-7}	0.84
Host-Normalized Offsets	2.6×10^{-4}	1.3×10^{-2}	...	5.2×10^{-5}	...	0.17	...	0.21
Fractional Fluxes	0.79	5.4×10^{-5}	3.0×10^{-2}	0.52	0.09	...

Note. — The values reported here for the physical offsets, host-normalized offsets, and fractional fluxes include only bursts with $\sigma_{R_{\text{phys}}} \lesssim 1.0$ kpc, $\sigma_{R_{\text{norm}}} \lesssim 0.5$, and error circle to galaxy area ratio $\lesssim 0.1$, respectively. Comparison samples of Type Ib/c and II SNe (physical offsets), Type Ib/c and II SNe (host-normalized offsets), Type Ic and II SNe (fractional fluxes), CCSNe, SLSNe, LGRBs - F06, and LGRBs - B02 are from Prieto et al. (2008), Kelly & Kirshner (2012), Kelly et al. (2008), Svensson et al. (2010), Lunnan et al. (2015), Fruchter et al. (2006), and Bloom et al. (2002), respectively.

^a For the host-normalized offsets, 80% of LGRB Monte Carlo synthetic distributions have p-values > 0.05 when compared to Type Ib/c SNe.

Table 5
Summary of LGRB Offset and Fractional Flux Medians

	Sample Median	Mean of Monte Carlo Medians	90% Confidence Interval
Physical Offsets	1.27 kpc	1.34 kpc	1.17 – 1.51 kpc
Host-Normalized Offsets	0.54	0.67	0.58 – 0.76
Fractional Fluxes	0.75

Note. — The values reported here for the physical offsets, host-normalized offsets, and fractional fluxes include only bursts with $\sigma_{R_{\text{phys}}} \lesssim 1.0$ kpc, $\sigma_{R_{\text{norm}}} \lesssim 0.5$, and error circle to galaxy area ratio $\lesssim 0.1$, respectively.